

THE WELL-POSEDNESS OF THE CAUCHY PROBLEM FOR A SYSTEM OF THERMOELASTICITY WITH SINGULAR COEFFICIENTS

AKBAR B. ALIEV AND GUNAY R. GADIROVA

Abstract. We consider the Cauchy problem for a system of thermoelasticity with some non-Lipschitz coefficients. We prove well-posedness of the corresponding Cauchy problem in some functional spaces.

1. Introduction and main results.

The various problems of thermoelasticity are reduced to the Cauchy problem or the mixed problem of hyperbolic-parabolic coupled systems. When coefficients are sufficiently smooth, these problems were studied by various authors [5], [8]-[20]. In the mentioned works the well-posedness of the corresponding Cauchy problem or the mixed problem, and the behavior of solutions, were investigated. In this paper, we consider the Cauchy problem:

$$\begin{cases} u_{tt} - a(t)\Delta u + \operatorname{div}\theta = 0 \\ \theta_t - \Delta\theta + \operatorname{div}u_t = 0 \end{cases} \quad t \in [0, T], \quad x \in R^n \quad (1.1)$$

with initial conditions

$$u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x), \quad \theta(0, x) = \theta_0(x). \quad (1.2)$$

It is known that if $a(t) \geq a_0 > 0$ satisfies the Lipschitz condition and $u_0 \in H^s(R^n)$, $u_1 \in H^{s-1}(R^n)$, $\theta_0 \in H^s(R^n)$, then problem (1.1)-(1.2) has a unique solution $u \in C([0, T]; H^s(R^n)) \cap C^1([0, T]; H^{s-1}(R^n))$, $\theta \in C([0, T], H^s(R^n))$ [1]-[4].

If we reject the Lipschitz condition, then this result, generally speaking is not valid. In this paper our interest is to investigate the case when $a(t)$ is not from the class C^1 at $t = 0$. Naturally, to investigate such problem there must be some condition on the singularity order of $a'(t)$ at $t \rightarrow 0$. But then it is necessary to understand the relation between this singular behavior and the right classes of well-posedness. Indeed, the results for the hyperbolic equation with singular coefficient in this direction for example were obtained in the works [1]-[4], [7]. Studies of hyperbolic-parabolic coupled system in this direction are few. In the article [6], the loss of smoothness of the solutions to the one class system of thermoelasticity with logarithmical Lipschitz coefficient is investigated. In this

2010 *Mathematics Subject Classification.* 34L10, 30E25, 34B24.

Key words and phrases. thermoelasticity, Cauchy problem, well-posedness, system of parabolic-hyperbolic equations.

paper we investigate the Cauchy problem for the system (1.1), (1.2) with singular coefficients $a(t)$.

At first we give some auxiliary notations. $\bigcap_{\beta>0} H^\beta(R^n)$ is denoted by H^∞ .

For $s \geq 1$ we'll denote by $\gamma_\beta^{(s)}$ the functional space with the norm $\|f\|_{\gamma_\beta^{(s)}} = \left\{ \int_{R^n} \exp(\beta|\xi|^{\frac{1}{s}}) |\hat{f}(\xi)|^2 d\xi \right\}^{\frac{1}{2}}$, where $\hat{f}(\xi)$ is the Fourier transformation of $f(x)$, i.e. $\hat{f}(\xi) = F[f](\xi)$

By $\gamma^{(s)}$ we denote $\gamma^{(s)} = \bigcap_{\beta>0} \gamma_\beta^{(s)}$.

Theorem 1.1. *Let*

$$a(t) \in C^1(0, T], \quad a(t) \geq a_0, \quad t \in (0, T], \quad (1.3)$$

$$t \cdot |a'(t)| \leq c, \quad t \in (0, T] \quad (1.4)$$

where $a_0, c \in (0, \infty)$, then Cauchy problem (1.1), (1.2) is well-posed in $H^\infty \times H^\infty \times H^\infty$.

Theorem 1.2. *Let (1.3) be satisfied. Suppose that there exist $c_1, c_2 > 0$ such that, for all $t \in [0, T]$*

$$t^q |a'(t)| \leq c_1, \quad (1.5)$$

$$t^p a(t) \leq c_2, \quad (1.6)$$

where $q > 1$ and $p \in [0, 1]$ with $p < q - 1$, then the Cauchy problem (1.1)-(1.2) is $\gamma^{(s)} \times \gamma^{(s)} \times \gamma^{(s)}$ well-posed for all $s < \frac{q-p}{q-1}$.

2. Proof of theorem 1.1.

The proof of theorem 1.1 and 1.2 is carried out by standard regularization method that based on some energetic estimation.

First of all, we observe that (1.4) implies that $a(\cdot) \in L_1(0, T)$, and there exist $c_1, c_2 > 0$ such that

$$|a(t)| \leq c_1 + c_2 \ln\left(1 + \frac{1}{t}\right), \quad (2.1)$$

We define for all $(t, \xi) \in [0, T] \times R^n \setminus \{0\}$

$$\tilde{a}(t, \xi) = \begin{cases} a(T) & \text{if } T|\xi| \leq 1, \\ a(|\xi|^{-1}) & \text{if } T|\xi| > 1 \text{ and } t|\xi| \leq 1, \\ a(t) & \text{if } t|\xi| > 1, \end{cases} \quad (2.2)$$

For all $(t, \xi) \in [0, T] \times R^n \setminus \{0\}$ we define also

$$d(t, \xi) = \begin{cases} |\tilde{a}(t, \xi) - a(t)| \cdot |\xi| & \text{if } t|\xi| \leq 1, \\ \frac{|a'(t)|}{a(t)} & \text{if } t|\xi| > 1, \end{cases} \quad (2.3)$$

and $k_\lambda(t, \xi) = (1 + |\xi|^2)^\lambda \exp(-\int_0^t d(s, \xi) ds)$.

We denote by $v(t, \xi)$ and $w(t, \xi)$ a Fourier transformation of $u(t, x)$ and $\theta(t, x)$, respectively with respect to the variable x , i.e. $v(t, \xi) = F[u](t, \xi)$, $w(t, \xi) = F[\theta](t, \xi)$. Let's define the weighted energetic function in the following way

$$E_\lambda(t) = \int_{R^n} E_\lambda(t, \xi) d\xi,$$

where

$$E_\lambda(t, \xi) = \left[|\dot{v}(t, \xi)|^2 + (1 + \tilde{a}(t, \xi) |\xi|^2) \cdot |v(t, \xi)|^2 + |w(t, \xi)|^2 \right] \cdot k_\lambda(t, \xi) \quad (2.4)$$

From (2.1) and (2.2) we deduced that for all $(t, \xi) \in [0, T] \times R^n \setminus \{0\}$

$$\tilde{a}(t, \xi) \leq c_3 + c_4 \ln(1 + |\xi|). \quad (2.5)$$

Lemma 2.1. *There exists $M > 0$, such that*

$$E_\lambda(t) \leq M E_\lambda(0), t \in [0, T].$$

Proof. After simple transformations we get

$$\begin{aligned} \frac{dE_\lambda(t, \xi)}{dt} &= [2Re\dot{v}(t, \xi) \cdot \bar{v}(t, \xi) + 2Re\dot{w}(t, \xi) \cdot \bar{w}(t, \xi) \\ &+ \dot{\tilde{a}}(t, \xi) |\xi|^2 \cdot |v(t, \xi)|^2 + \left(1 + \tilde{a}(t, \xi) |\xi|^2\right) 2 \cdot Re\dot{v}(t, \xi) \cdot \bar{v}(t, \xi)] \\ &\quad \times k(t, \xi) - E_\lambda(t, \xi) \cdot d(t, \xi). \end{aligned} \quad (2.6)$$

Since

$$Re \sum_{k=1}^n i \xi_k [w \cdot \dot{v} + \dot{w} \cdot \bar{v}] = 0,$$

the equalities (1.1) and (2.6) show that

$$\begin{aligned} \frac{dE_\lambda(t, \xi)}{dt} &= \left[[1 + (\tilde{a}(t, \xi) - a(t)) |\xi|^2] \cdot |v(t, \xi)|^2 \right. \\ &\quad \left. + \dot{\tilde{a}}(t, \xi) \cdot |\xi|^2 |v(t, \xi)|^2 - |\xi|^2 \cdot |w(t, \xi)|^2 \right] \cdot k(t, \xi) - E_\lambda(t, \xi) d(t, \xi). \end{aligned} \quad (2.7)$$

If $t|\xi| > 1$, by (2.2) and (2.3) we obtain that

$$\tilde{a}(t, \xi) - a(t) = 0, d(t, \xi) = \frac{|\dot{a}(t)|}{a(t)} \quad (2.8)$$

and hence

$$\dot{\tilde{a}}(t, \xi) |\xi|^2 k(t, \xi) - E_\lambda(t, \xi) d(t, \xi) \leq (\dot{a}(t) - |\dot{a}(t)|) |\xi|^2 k(t, \xi) \leq 0. \quad (2.9)$$

By virtue of (1.1), (2.8) and (2.9), it follows from (2.7) that

$$\frac{dE_\lambda(t, \xi)}{dt} \leq 0. \quad (2.10)$$

As $t|\xi| < 1$, then we have

$$2(\dot{\tilde{a}}(t, \xi) - a(t)) Re\dot{v}(t, \xi) \bar{v}(t, \xi) - (\dot{\tilde{a}}(t, \xi) - a(t)) [|\dot{v}(t, \xi)|^2 + |v(t, \xi)|^2] \leq 0.$$

Therefore

$$\frac{dE_\lambda(t, \xi)}{dt} \leq M \left[|\dot{v}(t, \xi)|^2 + |v(t, \xi)|^2 \right] \cdot k(t, \xi) \leq M E_\lambda(t, \xi). \quad (2.11)$$

By applying the Gronwall inequality, from (2.10), (2.11) we get $E_\lambda(t, \xi) \leq M_T E(0, \xi)$, where $M_T = e^{MT}$. Thus

$$E_\lambda(t) \leq M_T E(0) \quad (2.12)$$

□

Lemma 2.2. *For any $\lambda \geq 0$ and $\varepsilon > 0$*

$$E_\lambda(t) \leq \Phi_{\lambda+\varepsilon}(t), \quad t \in [0, T] \quad (2.13)$$

where

$$\Phi_\alpha(t) = \int_{R^n} (1 + |\xi|^2)^\alpha \left[|\dot{v}(t, \xi)|^2 + (1 + |\xi|^2) |v(t, \xi)|^2 + |w(t, \xi)|^2 \right] d\xi. \quad (2.14)$$

Proof. Inequality (2.13) is resulted from inequality (2.5). □

Lemma 2.3. *There exists $N > 0$ and $\Lambda > 0$ such that for all $t \in [0, T]$,*

$$E_\delta(t) \geq \Lambda \cdot \Phi_{\delta-N}(t).$$

Proof. Let $T|\xi| \leq 1$. By the definition of $d(s, \xi)$ we have

$$\int_0^t d(s, \xi) ds \leq \int_0^T |\tilde{a}(t, \xi) - a(t)| \cdot \frac{1}{T} dt \leq |\tilde{a}(t, \xi)| + \frac{1}{T} \|a(\cdot)\|_{L_1(0, T)}. \quad (2.15)$$

Secondly, we consider the case $T|\xi| > 1$. From (2.2) and (2.15) we have that

$$\begin{aligned} \int_0^t d(s, \xi) ds &\leq \int_0^T d(s, \xi) d\xi \leq \int_0^{|\xi|^{-1}} |\tilde{a}(t, \xi) - a(t)| \cdot |\xi| dt \\ &+ \int_{|\xi|^{-1}}^T \frac{|\dot{a}(t, \xi)|}{a(t)} dt \leq c_1 + c_2 \ln(1 + |\xi|) + |\xi| \int_0^{|\xi|^{-1}} a(t) dt \\ &+ \int_{|\xi|^{-1}}^T \frac{C}{t} dt \leq c_3 + c_4 \ln(1 + |\xi|). \end{aligned} \quad (2.16)$$

Hence, from (2.4) and (2.5) we get

$$e^{-\int_0^t d(s, \xi) ds} \geq c_5 (1 + |\xi|)^{-c_4}. \quad (2.17)$$

Hence there exist $N > 0$ and $c > 0$ such that for all $t \in [0, 1]$

$$E_\lambda(t) \geq c \Phi_{\lambda-N}(t)$$

From lemmas 2.1-2.3 it follows that the Cauchy problem (1.1), (1.2) is well-posed in $H^\infty \times H^\infty \times H^\infty$. □

3. Proof of theorem 1.2.

Let $1 < s < \frac{q-p}{q-1}$. At first we introduce some notation :

$$\tilde{a}_1(t, \xi) = \begin{cases} a(T) & \text{if } T |\xi|^{\frac{1}{s(q-1)}} \leq 1, \\ a(|\xi|^{-\frac{1}{s(q-1)}}) & \text{if } T |\xi|^{\frac{1}{s(q-1)}} > 1, t |\xi|^{\frac{1}{s(q-1)}} \leq 1, \\ a(t) & \text{if } t |\xi|^{\frac{1}{s(q-1)}} > 1, \end{cases}$$

$$d_1(t, \xi) = \begin{cases} |\tilde{a}(t, \xi) - a(t)| \cdot |\xi| & \text{if } t |\xi|^{\frac{1}{s(q-1)}} \leq 1, \\ \frac{|\tilde{a}(t, \xi)|}{a(t)} & \text{if } t |\xi|^{\frac{1}{s(q-1)}} > 1 \end{cases}$$

and

$$\phi_{\lambda, \beta}(t, \xi) = e^{-\int_0^t d_1(s, \xi) ds + \beta |\xi|^{\frac{1}{\lambda}}},$$

where β is a positive constant.

We also introduce the notation

$$H_{\lambda, \beta}(t) = \int_{R^n} H_{\lambda, \beta}(s, \xi) ds$$

where

$$H_{\lambda, \beta}(s, \xi) = \left[|\dot{v}(t, \xi)|^2 + (1 + \tilde{a}(t, \xi) |\xi|^2) |v(t, \xi)|^2 + |w(t, \xi)|^2 \right] \phi_{\lambda, \beta}(t, \xi).$$

Using (2.6) we get that

$$H_{\lambda, \beta}(t) \leq M \int_{R^n} e^{\beta' |\xi|^{\frac{1}{\lambda}}} \left[|\dot{v}(t, \xi)|^2 + (1 + |\xi|^2) (|v(t, \xi)|^2 + |w(t, \xi)|^2) \right] d\xi. \quad (3.1)$$

On the other hand

$$\int_0^t d_1(s, \xi) ds \leq \delta (|\xi|^{\frac{1}{\lambda}} + 1), i = 1, 2. \quad (3.2)$$

Thus, we have

$$\phi_{\lambda, \beta}(t, s) \geq e^{-2\delta + (\beta - 2\delta) |\xi|^{\frac{1}{\lambda}}}$$

and

$$H_{\lambda, \beta}(t) \geq e^{-2\delta} \int_{R^n} e^{\beta' |\xi|^{\frac{1}{\lambda}}} \left[|\dot{v}(t, \xi)|^2 + (1 + |\xi|^2) (|v(t, \xi)|^2 + |w(t, \xi)|^2) \right] d\xi, \quad (3.3)$$

where $\beta' = \beta - 2\delta$.

In the same way as lemma 2.1, we can show that there exists $M > 0$ such that

$$H_{\lambda, \beta}(t) \leq M H_{\lambda, \beta}(0), t \in [0, T]. \quad (3.4)$$

From (3.1)-(3.4) it follows that the Cauchy problem (1.1), (1.2) is well-posed in $\gamma^{(s)} \times \gamma^{(s)} \times \gamma^{(s)}$.

Acknowledgements

The authors thank the referee for his valuable comments on the paper.

References

- [1] A.B. Aliev, G.D. Shukurova, Well-posednes of the Cauchy problem for hyperbolic equations with non-lipschitz coefficients, *Abstract and Applied Analysis*, **2009**, Article ID 182371, 15 pages.
- [2] M. Cicognani, F. Colombini, Modulus of continuity of the coefficients and loss of derivatives in the strictly hyperbolic Cauchy problem, *Journal of Differential Equations*, **221** (1), (2006) 143-157.
- [3] F. Colombini, D. Del Santo, T. Kinoshita, Well-posedness of the Cauchy problem for hyperbolic equations with non Lipschitz coefficients, *Annali della Scuola Normale Superiore di Pisa*, **1**, (2002) 327-358.
- [4] F. Colombini, E.S. De Gorge, Existence et unique des solutions des equations hyperboliques du second orde a coefficients no dependant, *Comptes Rendus de l'Academie des Sciences*, **286**, (1978) 1045-1051.
- [5] C.M. Dafermos, On the existence and the asymptotic stability of solutions to the equations of linear thermoelasticit, *Arch. Rational Mech. Anal.*, **29**, (1968) 241-271
- [6] G.R. Gadirova, Loss of smoothness of solution of a hyperbolic-parabolic system with singular coefficient, *Transactions of NAS of Azerbaijan*, **XXXIV** (4), (2014) 37-44.
- [7] F. Hirose, Loss of regularity for second order hyperbolic equations with singular coefficients, *Osaka Journal of Mathematics*, **42**, (4), (2005) 767-790 .
- [8] W.J. Hrusa, M.A. Tarabek, On smooth solutions of the Cauchy problem in one-dimensional nonlinear thermoelasticity, *Quart. Appl. Math.*, **47** (1989), 631-644.
- [9] L. Hsiao, S. Jiang, Nonlinear hyperbolic-parabolic coupled systems, in Handbook of Diferential Equations, *Evolutionary Equations (Dafermos, C.M. & Feireisl, E. (Eds.)) Elsevier*, **1**, (2004) 287-384.
- [10] S. Jiang, Global smooth solutions to a one-dimensional nonlinear thermoviscoelastic model, *Adv.Math Sci Appl*, **7**, (1997), 771-787.
- [11] S. Jiang, Global existence of smooth solutions in one-dimensional nonlinear thermoelasticity, *Proc. R. Soc. Edinburgh*, **115A** (1990), 257-274.
- [12] S. Jiang, Global solutions of the Neumann problem in one-dimensional nonlinear thermoelasticity, *Nonlinear Anal., TMA*, **19** (1992), 107-121.
- [13] S. Jiang, J.E. Musimnoz Rivera, R. Racke, Asymptotic stability and global existence in thermoelasticity with symmetry, *Quart. Appl. Math.*, **56** (1998), 259-275.
- [14] G. Ponce, R. Racke, Global existence of small solutions to the initial value problem for nonlinear thermoelasticity, *J. Dif. Equat.*, **87** (1990), 70-83.
- [15] R. Racke, Y. Shibata, S. Zheng, Global solvability and exponential stability in one-dimensional nonlinear thermoelasticity, *Quart. Appl. Math.* **51** (1993), 751-763.
- [16] R. Racke, Y.G. Wang, Nonlinear well-posedness and rates of decay in thermoelasticity with second sound, *J. Hyperbolic Diferential Equations* **5** (2008), 25-43.
- [17] M. Reissig, About strictly hyperbolic operators with non-regular coefficients, *Pliska Studia, Mathematica Bulgarica*, **15**, (2003) 105-130.
- [18] Y.G. Wang, L. Yang, $L_p \rightarrow L_q$ decay estimates for Cauchy problems of linear thermoelastic systems with second sound in $3 - d$, *Proc. R. Soc. Edinburgh, Sec. A* **136**, (2006), 189-207
- [19] L. Yang, Y.G. Wang, Well-posedness and decay estimates for Cauchy problems of linear thermoelastic systems of type III in 3-D, *Indiana Univ. Math. J.*, **55** (2006), 1333-1362.
- [20] L. Yang, Y.G. Wang, Well-posedness and decay estimates for Cauchy problems of linear thermoelastic systems of type III in 3-D, *Indiana Univ Math J*, **4**, (2006), 1333-1364.

Azerbaijan Technical University, 25 Huseyn Cavid Ave., AZ1073, Baku, Azerbaijan

Institute of Mathematics and Mechanics, NAS of Azerbaijan, 9 B. Vahabzadeh str., AZ1141, Baku, Azerbaijan

E-mail address: alievakbar@gmail.com

Gunay R. Gadirova

Institute of Mathematics and Mechanics, NAS of Azerbaijan, 9 B. Vahabzadeh str., AZ1141, Baku, Azerbaijan

E-mail address: gunayqedirova@inbox.ru

Received: February 15, 2017; Revised: May 24, 2017; Accepted: May 31, 2017