

DIFFERENCE APPROXIMATION OF THE INVERSE PROBLEM OF DETERMINING THE HIGHEST COEFFICIENT IN A PARABOLIC EQUATION WITH INTEGRAL CONDITIONS

RAFIQ K. TAGIYEV AND SHAHLA I. MAHARRAMLI

Abstract. We consider the variational statement of the inverse problem of determining the higher coefficient of a parabolic equation with an integral boundary condition and an integral overdetermination condition. We establish estimates for the rate of convergence of difference approximations of the problem with respect to the state and functional.

1. Introduction

When we study variational statements of coefficient inverse problems for equations of mathematical physics, a number of difficulties arise due to their incorrectness, nonconvexity and describing the state of systems [1, 3]. These circumstances make it difficult to substantiate numerical methods, in particular. Variational statements of coefficient inverse problems for parabolic equations were studied in [1, 3, 4, 6, 7, 12] and others. Convergence of difference approximations of optimal control problems for the coefficients of parabolic equations and variational statements of coefficient inverse problems for these equations were studied in [8, 11] and others. However, these issues have been studied much less for parabolic equations with an integral overdetermination condition.

In this paper, we consider a variational statement of the inverse problem of determining the higher coefficient of a parabolic equation with an integral boundary condition and an integral overdetermination condition. We establish estimates for the rate of convergence of difference approximations of the problem with respect to the state and functional.

2. Variational statement of the inverse problem and its well-posedness

Suppose that it is required to find a pair of functions $\{v(x), u(x, t) = u(x, t; v)\}$ minimizing the functional

2010 *Mathematics Subject Classification.* 35K20, 49J20.

Key words and phrases. parabolic equation, inverse problem, integral condition, difference approximation.

$$J(v) = \int_0^l \left| \int_0^T \alpha(t) u(x, t; v) dt - \beta(x) \right|^2 dx \quad (2.1)$$

subject to the following boundary value problem

$$\frac{\partial u}{\partial t} - \frac{\partial}{\partial x} \left(v(x) \frac{\partial u}{\partial x} \right) + a(x, t) u = f(x, t), \quad (x, t) \in Q_T, \quad (2.2)$$

$$u(x, 0) = \varphi(x), \quad 0 \leq x \leq l, \quad (2.3)$$

$$u(0, t) = 0, \quad v(l) \frac{\partial u(l, t)}{\partial x} = \int_0^l H(x, t) u(x, t) dx, \quad 0 < t \leq T, \quad (2.4)$$

and the function $v(x)$ belongs to the set

$$V = \left\{ v = v(x) \in W_2^1(0, l) : 0 < \nu \leq v(x) \leq \mu, \right. \\ \left. |v'(x)| \leq \mu \text{ a.e. } (0, l) \right\}. \quad (2.5)$$

Here $l, T, \nu, \mu > 0$ are given numbers, $Q_T = \{(x, t) : 0 < x < l, 0 < t \leq T\}$, $a(x, t)$, $f(x, t)$, $\varphi(x)$, $H(x, t)$ are given measurable functions that satisfy the following conditions:

$$|a(x, t)| \leq \mu \text{ a. e. } Q_T, f(x, t) \in L_2(Q_T), \varphi(x) \in W_{2,0}^1(0, l), |H(x, t)| \leq \mu,$$

$$\left| \frac{\partial H(x, t)}{\partial t} \right| \leq \mu \text{ a.e. } Q_T,$$

$$\alpha(t) \in W_2^1(0, T), \beta(x) \in W_2^1(0, l). \quad (2.6)$$

In the work, the function spaces used and the norms correspond to [10, p.23]. In addition, positive constant which is depend on the estimated values and the steps of introduced grids, is denoted by M .

Direct and inverse problems with integral conditions have been studied by various authors in [2, 5, 9] and others.

By the solution of the boundary value problem (2.2)-(2.4) for each control $v = v(x) \in V$, we mean a generalized solution from $V_2^{1,0}(Q_T)$. Here $V_2^{1,0}(Q_T)$ is a subspace of $V_2(Q_T)$, whose elements have traces from $L_2(0, l)$ for all $t \in [0, T]$, continuously varying with $t \in [0, T]$ in the norm $L_2(0, l)$. It follows from the results of [14] that, under the above-mentioned assumptions (2.6), the boundary value problem (2.2)-(2.4) has a unique generalized solution from $V_2^{1,0}(Q_T)$ and this solution also belongs to the space $W_2^{1,1}(Q_T)$ for each fixed $v \in V$, and the following a priori estimate is valid

$$\|u\|_{V_2^{1,0}(Q_T)} + \max_{0 \leq t \leq T} \left\| \frac{\partial u}{\partial x}(x, t) \right\|_{2,(0,l)} + \left\| \frac{\partial u}{\partial t} \right\|_{2,Q_T} \leq \\ \leq M \left[\|\varphi\|_{2,(0,l)}^{(1)} + \|f\|_{2,Q_T} \right]. \quad (2.7)$$

In addition, it was proved in [14] that problem (2.1)-(2.5) is correctly posed in the weak topology of the space $W_2^1(0, l)$, i.e. the set of optimal control problems (2.1)-(2.5) $V_* = \{v_* = v_*(x) \in V : J(v_*) = J_* \equiv \inf \{J(v) : v \in V\}\}$ is nonempty and any minimizing sequence $\{v_n = v_n(x)\} \subset V$ of the functional (2.1) converges to the set V_* weakly in $W_2^1(0, l)$.

3. Difference approximation of the problem and its well-posedness

We introduce the following grids on the segments $[0, l]$ and $[0, T]$:

$$\begin{aligned}\omega_h &= \{x_i = ih \in [0, l] : i = 1, 2, \dots, N-1, h = l/N\}, \\ \omega_h^+ &= \{x_i = ih \in [0, l] : i = 1, 2, \dots, N\}, \\ \bar{\omega}_h &= \{x_i = ih \in [0, l] : i = 0, 1, \dots, N\}, \\ \omega_\tau &= \{t_j = j\tau \in [0, T] : j = 1, 2, \dots, L, \tau = T/L\}, \\ \bar{\omega}_{h*} &= \{\bar{x}_i = (i - 0.5)h \in [0, l] : i = 1, 2, \dots, N\}, \\ \omega_{h*}^0 &= \bar{\omega}_{h*} / \{\bar{x}_N = (N - 0.5)h\}.\end{aligned}$$

We introduce the following grids on the rectangle \bar{Q}_T :

$$\omega_T = \omega_h \times \omega_\tau, \omega_T^+ = \omega_h^+ \times \omega_\tau, \bar{\omega}_T = \bar{\omega}_h \times \omega_\tau, \omega_t^+ = \omega_h^+ \times \{t' = \tau, 2\tau, \dots, t\}.$$

Let $\bar{h} = \bar{h}(x) = h$, if $x \in \omega_h$, $\bar{h}(0) = \bar{h}(l) = 0.5h$.

We introduce the following scalar products and norms for grid functions defined on the corresponding grids:

$$\begin{aligned}(y, z)_{\omega_h^+} &= \sum_{\omega_h^+} \bar{h}yz, \|y\|_{2, \omega_h^+} = (y, y)_{\omega_h^+}^{1/2}, \\ (y, z)_{\omega_\tau} &= \sum_{\omega_\tau} \tau yz, \|y\|_{2, \omega_\tau} = (y, y)_{\omega_\tau}^{1/2}, (y_{\bar{x}}, z_{\bar{x}})_{\omega_h^+} = \sum_{\omega_h^+} h y_{\bar{x}} z_{\bar{x}}, \\ \|y_{\bar{x}}\|_{2, \omega_h^+} &= (y_{\bar{x}}, y_{\bar{x}})_{\omega_h^+}^{1/2}, (y, z)_{\omega_T^+} = \sum_{\omega_\tau} \tau (y, z)_{\omega_h^+}, \\ \|y\|_{2, \omega_T^+} &= (y, y)_{\omega_T^+}^{1/2}, (y_{\bar{x}}, z_{\bar{x}})_{\omega_t^+} = \sum_{t=\tau}^t \tau (y_{\bar{x}}, z_{\bar{x}})_{\omega_h^+}, \\ \|y_{\bar{x}}\|_{2, \omega_t^+} &= (y_{\bar{x}}, y_{\bar{x}})_{\omega_t^+}^{1/2}, t \in \omega_\tau, |y|_{\omega_t^+} \equiv \|y\|_{V_2^{1,0}(\omega_t^+)} = \\ &= \max_{0 \leq t' \leq t} \|y(x, t')\|_{2, \omega_h^+} + \|y_{\bar{x}}\|_{2, \omega_t^+}, \quad t \in \omega_\tau, (y_{\bar{t}}, z_{\bar{t}})_{\omega_t^+} = \sum_{t=\tau}^t \tau (y_{\bar{t}}, z_{\bar{t}})_{\omega_h^+}, \\ \|y_{\bar{t}}\|_{2, \omega_t^+} &= (y_{\bar{t}}, y_{\bar{t}})_{\omega_t^+}^{1/2}, t \in \omega_\tau, \\ \theta(t, y) &\equiv |y|_{\omega_t^+} + \sqrt{\tau} \|y_{\bar{t}}(x, t)\|_{2, \omega_t^+}, \quad t \in \omega_\tau, \\ \|v_h\|_{2, \bar{\omega}_{h*}} &= \left(\sum_{\bar{\omega}_{h*}} h v_h^2(x) \right)^{1/2}, \|v_h\|_{2, \bar{\omega}_{h*}}^{(1)} = \left[\sum_{\bar{\omega}_{h*}} h v_h^2(x) + \sum_{\omega_{h*}^0} h v_{hx}^2(x) \right]^{1/2}, \\ \|v_h\|_{\infty, \bar{\omega}_{h*}} &= \max_{x \in \bar{\omega}_{h*}} |v_h(x)|.\end{aligned}$$

We also introduce elementary cells:

$$e_1(x) = \{\xi : x - 0.5h \leq \xi < x + 0.5h\}, x \in \omega_h, e_1(0) = \{\xi : 0 \leq \xi < 0.5h\},$$

$$e_1(l) = \{\xi : l - 0.5h \leq \xi \leq l\}; e_2(t) = \{\theta : t - \tau \leq \xi < t\}, t \in \omega_\tau,$$

$$e_1^*(x) = \{\xi : x - h \leq \xi < x\}, x \in \omega_h^+; e(x, t) = e_1(x) \times e_2(t), (x, t) \in \bar{\omega}_T,$$

$$e^*(x, t) = e_1^*(x) \times e_2(t), \quad (x, t) \in \omega_T^+.$$

Let S^x , S^x , S_-^t be Steklov's one-dimensional averaging operators. In addition, let $S^{tx} = S_-^t S^x$ be the product of the averaging operators S^x and S_-^t .

We state the problem (2.1)-(2.5) in the following difference approximation form: it is required to find a pair of grid functions $\{v_h(x), y(x, t) = y(x, t; v_h)\}$ minimizing the grid functional

$$J_{h\tau}(v_h) = \sum_{x \in \omega_h^+} \bar{h} \left| \tau \sum_{t \in \omega_\tau} \alpha_\tau(t) y(x, t; v_h) - \beta_h(x) \right|^2 \quad (3.1)$$

subject to the difference boundary value problem

$$y_{\bar{t}} - (v_h(x - 0.5h) y_{\bar{x}})_x + a_{h\tau}(x, t) y = f_{h\tau}(x, t), \quad (x, t) \in \omega_T, \quad (3.2)$$

$$y(x, 0) = \varphi_h(x), \quad x \in \bar{\omega}_h, \quad (3.3)$$

$$y(0, t) = 0, v_h(l - 0.5h) y_{\bar{x}}(l, t) = \sum_{x \in \omega_h^+} h H_{h\tau}(x, t) y(x, t) - 0.5h [y_{\bar{t}}(l, t) + a_{h\tau}(l, t) y(l, t) - f_{h\tau}(l, t)], t \in \omega_\tau \quad (3.4)$$

and grid controls $v_h(x)$ are such that

$$v_h(x) \in V_h = \{v_h(x) \in W_2^1(\bar{\omega}_{h*}) :$$

$$0 < \nu \leq v_h(x) \leq \mu, x \in \bar{\omega}_{h*}, |v_{hx}(x)| \leq \mu, x \in \omega_{h*}^0 \}. \quad (3.5)$$

Here $a_{h\tau}(x, t) = S^{tx} a$, $f_{h\tau}(x, t) = S^{tx} f$, $H_{h\tau}(x, t) = S_-^x H$, $(x, t) \in \omega_T^+$, $\varphi_h(x) = S^x \varphi$, $\varphi_h(0) = 0$, $\beta_h(x) = S^x \beta$, $x \in \omega_h^+$; $\alpha_\tau(t) = S_-^t \alpha$, $t \in \omega_\tau$.

We represent the difference boundary value problem (3.2)-(3.4) in the following form:

$$y_{\bar{t}} = Ay + f_{h\tau}(x, t), \quad (x, t) \in \omega_T^+, \quad (3.6)$$

$$y(x, 0) = \varphi_h(x), \quad x \in \bar{\omega}_h, \quad (3.7)$$

$$y(0, t) = 0, \quad t \in \omega_\tau, \quad (3.8)$$

where

$$Ay = \begin{cases} (v_h(x - 0.5h) y_{\bar{x}})_x - a_{h\tau}(x, t) y, & (x, t) \in \omega_T, \\ -\frac{2}{h} v_h(l - 0.5h) y_{\bar{x}}(l, t) - a_{h\tau}(l, t) y(l, t) + \frac{2}{h} \sum_{\omega_h^+} h H_{h\tau}(x, t) y(x, t), & t \in \omega_\tau. \end{cases}$$

Theorem 3.1. *Let conditions (2.6) be satisfied and the step τ in the variable t satisfies the inequality:*

$$\tau < \tau_0 = \frac{1}{2} \left[2\mu + \mu^2 l + \frac{1}{\nu} + \frac{1}{l} \right]^{-1} \left(2 + \nu^{-1/2} \right)^{-2}. \quad (3.9)$$

Then problem (3.2)-(3.4) is uniquely solvable for each $v_h \in V_h$ and a priori estimate is valid

$$\begin{aligned} \|y(x, t; v_h)\|_{V_2^{1,0}(\omega_T^+)} + \sqrt{\tau} \|y_{\bar{t}}(x, t; v_h)\|_{2, \omega_T^+} &\leq \\ &\leq M \left[\|\varphi_h\|_{2, \omega_h^+} + 2 \|f_{h\tau}\|_{2,1, \omega_T^+} \right]. \end{aligned} \quad (3.10)$$

Proof. Let us carry out the main points of the proof of estimate (3.10). We multiply equation (3.6) and $\tau y(x, t')$ as a scalar product $(\cdot)_{\omega_h^+}$ and sum the resulting equality over t' from $t' = \tau$ to $t' = t$, where $t \in \omega_\tau$ is some point. Then using the summation by parts formulas and identity $y_{\bar{t}} y = 0.5(y^2)_{\bar{t}} + 0.5\tau y_{\bar{t}}^2$, $(x, t) \in \omega_\tau^+$ to equality

$$\begin{aligned} & \frac{1}{2} \|y(x, t)\|_{2, \omega_h^+}^2 + \tau \sum_{t'=\tau}^t h \sum_{x \in \omega_h^+} v_h(x - 0.5h) y_{\bar{x}}^2(x, t') + \\ & + \frac{1}{2} \tau^2 \sum_{t'=\tau}^t \|y_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 = \frac{1}{2} \|y(x, 0)\|_{2, \omega_h^+}^2 - \tau \sum_{t'=\tau}^t (a_{h\tau}(x, t'), \\ & y^2(x, t'))_{2, \omega_h^+} + \tau \sum_{t'=\tau}^t (H_{h\tau}(x, t'), y(x, t'))_{2, \omega_h^+} y(l, t') + \\ & + \tau \sum_{t'=\tau}^t (f_{h\tau}(x, t'), y(x, t'))_{2, \omega_h^+}. \end{aligned} \quad (3.11)$$

Then, using conditions (2.6), the Cauchy inequality with ε from [10, p.33]

$$|ab| \leq \frac{\varepsilon}{2} |a|^2 + \frac{1}{2\varepsilon} |b|^2, \quad \varepsilon > 0$$

for $\varepsilon = \mu\sqrt{l}$ and the inequality [13, p.290]

$$y^2(l, t) \leq \varepsilon \|y_{\bar{x}}(x, t)\|_{2, \omega_h^+}^2 + \left(\frac{1}{\varepsilon} + \frac{1}{l}\right) \|y(x, t)\|_{2, \omega_h^+}^2, \quad (3.12)$$

for $\varepsilon = \nu$, and majorizing the left and right sides of equality (3.11), we obtain the inequality

$$\begin{aligned} & \|y(x, t)\|_{2, \omega_h^+}^2 + \nu\tau \sum_{t'=\tau}^t \|y_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 + \tau^2 \sum_{t'=\tau}^t \|y_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 \leq \\ & \leq \gamma(t) \left[\|y(x, 0)\|_{2, \omega_h^+}^2 + 2\tau \sum_{t'=\tau}^t \|f_{h\tau}(x, t')\|_{2, \omega_h^+} \right] + \\ & + ct\gamma^2(t) \equiv \delta(t), \quad t \in \omega_\tau, \end{aligned} \quad (3.13)$$

where $\gamma(t) = \max_{0 \leq t' \leq t} \|y(x, t')\|_{2, \omega_h^+}$, $c = 2\mu + \mu^2 l + \nu^{-1} + l^{-1}$. From this inequality, we extract three consequences $\forall t \in \omega_\tau$

$$\begin{aligned} & \max_{0 \leq t' \leq t} \|y(x, t')\|_{2, \omega_h^+}^2 \leq \delta(t), \tau \sum_{t'=\tau}^t \|y_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 \leq \\ & \leq \nu^{-1} \delta(t), \tau \sum_{t'=\tau}^t \tau \|y_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 \leq \delta(t). \end{aligned}$$

We extract the square root from both parts of these inequalities, and add the resulting inequalities. Then, using condition (3.9), for $t < 2\tau_0$ we obtain the estimate

$$\begin{aligned} \theta(t, y) &\equiv \max_{0 \leq t' \leq t} \|y(x, t')\|_{2, \omega_h^+} + \left(\tau \sum_{t'=\tau}^t \|y_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} + \\ &+ \sqrt{\tau} \left(\tau \sum_{t'=\tau}^t \|y_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} \leq \left[1 - \left(2 + \nu^{-1/2} \right) \sqrt{ct} \right]^{-2} \left(2 + \nu^{-1/2} \right)^2 \times \\ &\times \left[\|y(x, 0)\|_{2, \omega_h^+} + 2\tau \sum_{t'=\tau}^t \|f_{h\tau}(x, t')\|_{2, \omega_h^+}^2 \right], \quad t \in \omega_\tau, \quad t < 2\tau_0. \end{aligned} \quad (3.14)$$

The interval $[0, T]$ is divided into subintervals $\Delta_1 = [0, \tau_0]$, $\Delta_2 = [\tau_0, 2\tau_0]$, \dots , Δ_n of a length not greater than τ_0 . Here $\Delta_n = [(n-1)\tau_0, T]$. Each of them satisfies the estimate (3.14). From these estimates, taking into account $\|y(x, t)\|_{2, \omega_h^+} \leq |y|_{\omega_t^+}$, $t \in \omega_\tau$, we derive inequality (3.10) for $t = T$. The unique solvability of problem (3.2)-(3.4) for each $v_h \in V_h$ is obvious. Theorem 3.1 is proved. \square

4. A priori estimate of the error of the difference method by state

Take arbitrary controls $v(\xi) \in V$, $v_h(x) \in V_h$. Let $u(\xi, \theta) = u(\xi, \theta; v)$ be a solution to the boundary value problem (2.2)-(2.4), and $y(x, t) = y(x, t; v_h)$ be a solution to the difference boundary value problem (3.2)-(3.4) corresponding to the controls $v \in V$ and $v_h(x) \in V_h$. Denote $\bar{u}(x, t) = \bar{u}(x, t; v)$ by the averaging of the function $u(\xi, \theta) = u(\xi, \theta; v)$, which is determined by the formula

$$\bar{u}(x, t) = \bar{u}(x, t; v) = \begin{cases} S_{-}^t u, & (x, t) \in \omega_T^+, \\ S^x \varphi(x), & x \in \bar{\omega}_h, \quad t = 0, \\ 0, & x = 0, \quad t \in \omega_\tau. \end{cases} \quad (4.1)$$

We will compare the grid solution $y(x, t; v_h)$ with the averaging $\bar{u}(x, t; v)$. Let $z = z(x, t) = z(x, t; v, v_h) = y(x, t; v_h) - \bar{u}(x, t; v)$, $(x, t) \in \bar{\omega}_T$ be the error of the state difference method. Using conditions (3.6)-(3.8) for the function z , we obtain the problem

$$z_{\bar{t}} = Az + \psi_{h\tau}(x, t), \quad (x, t) \in \omega_T^+, \quad (4.2)$$

$$z(x, 0) = 0, \quad x \in \bar{\omega}_h, \quad (4.3)$$

$$z(0, t) = 0, \quad t \in \omega_\tau, \quad (4.4)$$

where $\psi_{h\tau}(x, t) = f_{h\tau}(x, t) + A\bar{u} - \bar{u}_{\bar{t}}$, $(x, t) \in \omega_T^+$.

We will assume that the generalized solution $u(\xi, \theta) = u(\xi, \theta; v)$ from $V_2^{1,0}(Q_T)$ also belongs to the space $W_2^{2,1}(Q_T)$. Then, applying the averaging operator S^{tx}

to equation (2.2) at the nodes $(x, t) \in \bar{\omega}_T$, after some transformations we obtain the following representation for $\psi_{h\tau}$:

$$\psi_{h\tau}(x, t) = \begin{cases} \sum_{k=1}^2 \eta_{1x}^{(k)}(x, t) + \eta_2(x, t) + \eta_{3\bar{t}}(x, t), & (x, t) \in \omega_T, \\ -\frac{2}{h} \sum_{k=1}^2 \eta_1^{(k)}(x, t) + \eta_2(x, t) + \eta_{3\bar{t}}(x, t) + \frac{2}{h} \eta_4(t), & x = l, t \in \omega_\tau, \end{cases} \quad (4.5)$$

where

$$\eta_1^{(1)}(x, t) = v(x - 0.5h, t) \left[\bar{u}_{\bar{x}}(x, t) - S_-^t \frac{\partial u(x - 0.5h, t)}{\partial x} \right], \quad (4.6)$$

$$\eta_1^{(2)}(x, t) = [v_h(x - 0.5h) - v(x - 0.5h)] \bar{u}_{\bar{x}}(x, t), \quad (4.7)$$

$$\eta_2(x, t) = S^{tx}(a(x, t)u(x, t)) - S^{tx}(a(x, t))\bar{u}, \quad (4.8)$$

$$\eta_3(x, t) = S^x u(x, t) - \bar{u}(x, t), \quad (x, t) \in \omega_T^+, \quad (4.9)$$

$$\eta_4(t) = \sum_{x \in \omega_h^+} h S_-^x (H(x, t)) \bar{u}(x, t) - S_-^t \left(\int_0^l H(\xi, t) u(\xi, t) d\xi \right), \quad t \in \omega_\tau. \quad (4.10)$$

Theorem 4.1. *Let the conditions of theorem 3.1 be satisfied and the solution of problem (2.2)-(2.4) belong to the space $W_2^{2,1}(Q_T)$. Then, for the solution z of problem (4.2)-(4.4), the estimate*

$$\begin{aligned} & \|z(x, t)\|_{V_2^{1,0}(\omega_T^+)} + \sqrt{\tau} \|z_{\bar{t}}(x, t)\|_{2, \omega_T^+} \leq \\ & \leq M \left[\sum_{k=1}^2 \left\| \eta_1^{(k)} \right\|_{2, \omega_T^+} + \|\eta_2\|_{2, \omega_T^+} + \left(\sum_{t \in \omega_\tau} \|\eta_3(x, t)\|_{2, \omega_h^+}^2 \right)^{1/2} + \|\eta_4\|_{2, \omega_\tau} \right] \end{aligned} \quad (4.11)$$

holds true.

Proof. Multiplying equations (4.2) scalarly $(\cdot, \cdot)_{\omega_h^+}$ by $\tau z(x, t')$ and calculating similarly to the obtained equality (3.11), we have

$$\begin{aligned} & \frac{1}{2} \|z(x, t')\|_{2, \omega_h^+}^2 + \tau \sum_{t'=\tau}^t h \sum_{x \in \omega_h^+} v_h(x - 0.5h) z_{\bar{x}}^2(x, t') + \frac{1}{2} \tau^2 \sum_{t'=\tau}^t \|z_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 = \\ & = - \sum_{t'=\tau}^t (a_{h\tau}(x, t'), z^2(x, t'))_{2, \omega_h^+} - \tau \sum_{t'=\tau}^t \left(\sum_{k=1}^2 \eta_1^{(k)}(x, t'), z_{\bar{x}}(x, t') \right)_{2, \omega_h^+} + \\ & \quad + \tau \sum_{t'=\tau}^t (\eta_2(x, t'), z(x, t'))_{2, \omega_h^+} - \\ & \quad - \tau \sum_{t'=0}^{t-\tau} (\eta_3(x, t'), z_t(x, t'))_{2, \omega_h^+} + (\eta_3(x, t), z(x, t))_{2, \omega_h^+} + \\ & \quad + \tau \sum_{t'=\tau}^t \left[\sum_{\omega_h^+} h H_{h\tau}(x, t') z(x, t') + \eta_4(t') \right] z(l, t'). \end{aligned} \quad (4.12)$$

Using the method of item 2 for obtaining estimates of type (3.10), as well as the formula of summation by parts, inequalities (3.12), (3.14) can be obtained from (4.12) for any $t \in \omega_\tau$ of the following three estimates

$$\begin{aligned} \max_{0 \leq t' \leq t} \|z(x, t')\|_{2, \omega_h^+}^2 &\leq \delta(t), \tau \sum_{t'=\tau}^t \|z_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 \leq \nu^{-1} \delta(t), \\ \tau \sum_{t'=\tau}^t \tau \|z_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 &\leq \delta(t), \end{aligned} \quad (4.13)$$

where

$$\begin{aligned} \delta(t) &= c_1 t \max_{0 \leq t' \leq t} \|z(x, t')\|_{2, \omega_h^+} + 2c_2 \max_{0 \leq t' \leq t} \|z(x, t')\|_{2, \omega_h^+} \times \\ &\quad \times \left[\left(\tau \sum_{t'=\tau}^t \|\eta_2(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} + \|\eta^{(3)}(x, t)\|_{2, \omega_h^+} \right] + \\ + 2 \sum_{k=1}^2 &\left(\tau \sum_{t'=\tau}^t \|\eta_1^{(k)}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} \left(\tau \sum_{t'=\tau}^t \|z_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} + \\ + 2 &\left(\tau \sum_{t'=0}^{t-\tau} \|\eta_3(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} \left(\tau \sum_{t'=\tau}^t \tau \|z_{\bar{t}}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} + \\ + 2c_3 &\left(\tau \sum_{t'=\tau}^t \eta_4^2(t') \right)^{1/2} \left[\left(\tau \sum_{t'=\tau}^t \|z_{\bar{x}}(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} + \right. \\ &\quad \left. + \left(\tau \sum_{t'=\tau}^t \|z(x, t')\|_{2, \omega_h^+}^2 \right)^{1/2} \right]. \end{aligned}$$

Here $c_1 = 2\mu + \mu^2 l + \frac{1}{\nu} + \frac{1}{l}$, $c_2 = \max(\sqrt{T}; 1)$, $c_3 = (\max(\nu; \frac{1}{\nu} + \frac{1}{l}))^{1/2}$.

Extracting the square root from both parts of inequality (4.13), adding the resulting inequality, after some transformations we arrive at estimate (4.11). Theorem 4.1 is proved. \square

Lemma 4.1. *Let the conditions of Theorem 3.1 be satisfied. Then the approximation error components (4.6)-(4.10) satisfy the estimates*

$$\|\eta_1^{(1)}\|_{2, \omega_T^+} \leq h \|v\|_{\infty, (0, l)} \left\| \frac{\partial^2 u}{\partial \xi^2} \right\|_{2, Q_T}, \quad (4.14)$$

$$\|\eta_1^{(2)}\|_{2, \omega_T^+} \leq \|v_h(x) - v(x)\|_{\infty, \bar{\omega}_{h^*}} \left\| \frac{\partial u}{\partial \xi} \right\|_{2, Q_T}, \quad (4.15)$$

$$\|\eta_2\|_{2, \omega_T^+} \leq \sqrt{2} \|a\|_{\infty, Q_T} \left[3h \left\| \frac{\partial u}{\partial \xi} \right\|_{2, Q_T} + \tau \left\| \frac{\partial u}{\partial \theta} \right\|_{2, Q_T} \right], \quad (4.16)$$

$$\left(\sum_{t \in \omega_\tau} \|\eta^{(3)}(x, t)\|_{2, \omega_h^+}^2 \right)^{1/2} \leq M \left(\frac{h^{3/2}}{\tau^{1/2}} + \tau^{1/2} \right) \|u\|_{2, Q_T}^{(2,1)}, \quad (4.17)$$

$$\|\eta_4\|_{2,\omega_\tau} \leq \sqrt{2l} \|H\|_{\infty,Q_T}^{(0,1)} \left[h^{3/2} \left\| \frac{\partial u}{\partial \xi} \right\|_{2,Q_T} + h^{1/2} \tau \|u\|_{2,Q_T} \right]. \quad (4.18)$$

Proof. Let us consider only the proof of estimates (4.18).

Using (4.1) and (4.10) it can be shown that the function $\eta_4(t)$, $t \in \omega_\tau$ has the representation

$$\begin{aligned} \eta_4(t) &= \frac{1}{\tau} \int_{t-\tau}^t \left[\sum_{x \in \omega_h^+} \int_{x-h}^x \left(H(\xi, t) \int_{\xi}^x \frac{\partial u(\xi_1, \theta)}{\partial \xi_1} d\xi_1 \right) d\xi \right] d\theta + \\ &+ \frac{1}{\tau} \int_{t-\tau}^t \left[\sum_{x \in \omega_h^+} \int_{x-h}^x \left(u(\xi, \theta) \int_{\theta}^t \frac{\partial H(\xi, \theta_1)}{\partial \theta_1} d\theta_1 \right) d\xi \right] d\theta. \end{aligned}$$

Evaluating this representation, we get

$$\begin{aligned} |\eta_4(t)| &\leq \frac{h^{3/2}}{\tau^{1/2}} \sum_{x \in \omega_h^+} \max_{x-h \leq \xi \leq x} |H(\xi, t)| \left\| \frac{\partial u}{\partial \xi} \right\|_{2,e^*(x,t)} + \\ &+ h^{1/2} \tau^{1/2} \sum_{x \in \omega_h^+} \max_{(\xi, \theta) \in e^*(x,t)} \left| \frac{\partial H(\xi, \theta)}{\partial \theta} \right| \|u\|_{2,e^*(x,t)}, \quad t \in \omega_\tau. \end{aligned}$$

Based on the resulting inequality, we establish estimate (4.18).

Estimates (4.14), (4.15), (4.16) are proved in a similar way. Estimate (4.17) was proved in [11]. \square

Theorem 4.2. *Let the conditions of Theorem 4.1 be satisfied. Then for any controls $v \in V$ and $v_h \in V_h$ the estimate of the error of the difference method with respect to the state is*

$$\begin{aligned} &\|y(x, t; v_h) - \bar{u}(x, t; v)\|_{V_2^{1,0}(\omega_T^+)} + \sqrt{\tau} \|y_{\bar{t}}(x, t; v_h) - \bar{u}_{\bar{t}}(x, t; v)\|_{2,\omega_T^+} \leq \\ &\leq M \left[h + \frac{h^{3/2}}{\tau^{1/2}} + \tau^{1/2} + \|v_h(x) - v(x)\|_{\infty, \bar{\omega}_{h^*}} \right] = ME(h, \tau, v, v_h). \quad (4.19) \end{aligned}$$

Corollary 4.1. *We can obtain various estimates of the convergence rate of the difference method with respect to the state from (4.19). Let $v_h(x) = v(x)$, $x \in \bar{\omega}_{h^*}$. Then the difference boundary value problem (3.2)-(3.4) has convergence rate estimates $O(h^{1/2})$ for $\tau \sim h$ or $\tau \sim h^2$ in the grid norm $V_2^{1,0}(\omega_T^+)$, convergence rate estimates $O(h^{5/8})$ for $\tau \sim h^{5/4}$ or $\tau \sim h^{7/4}$, convergence rate estimates $O(h^{3/4})$ for $\tau \sim h^{3/2}$.*

5. Estimates of the error and convergence rate of approximations with respect to the functional

Theorem 5.1. *Let the conditions of Theorem 4.1 be satisfied. Then, for any controls $v \in V$ and $v_h \in V_h$ for the error of the grid functional (3.2), we have the estimate*

$$|J(v) - J_{h\tau}(v_h)| \leq M [h + \tau + E(h, \tau, v, v_h)]. \quad (5.1)$$

To study the relationship between problems (2.1)-(2.5) and (3.1)-(3.5), we introduce two mappings $P_h : W_2^1(\omega_h^+) \rightarrow W_2^1(0, l)$ and $Q_h : W_2^1(0, l) \rightarrow W_2^1(\omega_h^+)$, acting according to the rules:

$$P_h v_h(\xi) = \begin{cases} v_h(0.5h), & 0 \leq \xi \leq 0.5h, \\ v_h(x - 0.5h) + v_{hx}(x - 0.5h)(\xi - x + 0.5h), & x - 0.5h \leq \xi \leq x + 0.5h, x \in \omega_h, \\ v_h(l - 0.5h), & l - 0.5h \leq \xi \leq l, \end{cases}$$

$$Q_h v(x) = v(x - 0.5h), \quad x \in \omega_h^+,$$

It is easy to show that for any controls $v_h \in V_h$, $v \in V$, there are inclusion places $P_h v_h(\xi) \in V$, $Q_h v(x) \in V_h$.

Lemma 5.1. *For arbitrary controls $v \in V$, $v_h \in V_h$, we have the estimates*

$$\max \{ |J(v) - J_{h\tau}(Q_h v)|, |J(P_h v_h) - J_{h\tau}(v_h)| \} \leq M \left[h + \frac{h^{3/2}}{\tau^{1/2}} + \tau^{1/2} \right].$$

These estimates follow from (5.1) and are based on the definitions of mappings P_h , Q_h .

Theorem 5.2. *Let the conditions of Theorem 4.1 be satisfied. Then approximations (3.1)-(3.5) satisfy the estimate*

$$|J_{h\tau^*} - J_*| \leq M \left[h + \frac{h^{3/2}}{\tau^{1/2}} + \tau^{1/2} \right]. \quad (5.2)$$

Here $J_{h\tau^*} \equiv \inf \{ J_{h\tau}(v_h) : v_h \in V_h \}$.

If, in addition, the sequence of controls $\{v_{h\varepsilon}\}$ is determined from the conditions

$$v_{h\tau} \in V_h, J_{h\tau^*} \leq J_{h\tau}(v_{h\varepsilon}) \leq J_{h\tau^*} + \varepsilon_{h\tau}, \quad \varepsilon_{h\tau} \geq 0, \quad \varepsilon_{h\tau} \rightarrow 0 \text{ for } h, \tau \rightarrow 0 \quad (5.3)$$

then the sequence $\{P_h v_{h\varepsilon}\}$ satisfies the estimate

$$0 \leq J(P_h v_{h\varepsilon}) - J_* \leq M \left[h + \frac{h^{3/2}}{\tau^{1/2}} + \tau^{1/2} \right] + \varepsilon_{h\tau}. \quad (5.4)$$

Proof. Let's take some control $v_* \in V_*$. It's obvious that $Q_h v_* \in V_h$. From this and Lemma 4.1 it follows

$$J_{h\tau^*} \leq J_{h\tau}(Q_h v_*) \leq J(v_*) + M \left[h + \tau^{1/2} + h^{3/2}\tau^{-1/2} \right] = J_{h*} + M \left[h + \tau^{1/2} + h^{3/2}\tau^{-1/2} \right]. \quad (5.5)$$

Let's take some control

$$v_{h*} \in V_{h*} = \{v_{h*} \in V_h : J_{h\tau}(v_{h*}) = J_{h\tau^*} \equiv \inf \{ J_{h\tau}(v_h) : v_h \in V_h \} \}.$$

Then it is clear that $P_h v_{h*} \in V_h$. From this and Lemma 4.1 it follows

$$\begin{aligned} J_* &\leq J(P_h v_{h*}) \leq J_{h\tau}(v_{h*}) + M \left[h + \tau^{1/2} + h^{3/2}\tau^{-1/2} \right] = \\ &= J_{h\tau^*} + M \left[h + \tau^{1/2} + h^{3/2}\tau^{-1/2} \right]. \end{aligned} \quad (5.6)$$

Estimates (5.2) follow from (5.5), (5.6).

Consider the sequence $\{v_{h\varepsilon}\} \subset V_h$ from (5.3). Then $P_h v_{h\varepsilon} \in V$ and

$$0 \leq J(P_h v_{h\varepsilon}) - J_* = [J(P_h v_{h\varepsilon}) - J_{h\tau}(v_{h\varepsilon})] + [J_{h\tau}(v_{h\varepsilon}) - J_{h\tau*}] + [J_{h\tau*} - J_*].$$

From this, and also from inequalities (5.2), (5.3) and Lemma 4.1 it follows estimate (5.4). Theorem 5.2 is proved. \square

Corollary 5.1. *From (5.2) and (5.4) one can obtain different estimates of the rate of convergence. For example, for $\tau \sim h$ or $\tau \sim h^2$ the estimates $|J_{h\tau*} - J_*| \leq Mh^{1/2}$, $0 \leq J_{h\tau}(v_{h\varepsilon}) - J_* \leq Mh^{1/2} + \varepsilon_{h\tau}$, for $\tau \sim h^{5/4}$ or $\tau \sim h^{7/4}$ the estimates $|J_{h\tau*} - J_*| \leq Mh^{5/8}$, $0 \leq J_{h\tau}(v_{h\varepsilon}) - J_* \leq Mh^{5/8} + \varepsilon_{h\tau}$, for $\tau \sim h^{3/2}$ the estimates $|J_{h\tau*} - J_*| \leq Mh^{3/4}$, $0 \leq J_{h\tau}(v_{h\varepsilon}) - J_* \leq Mh^{3/4} + \varepsilon_{h\tau}$ are valid.*

Moreover, in these cases the sequence of controls $\{P_h v_{h\varepsilon}\}$ is minimizing for problem (2.1)-(2.5) and converges $W_2^1(0, l)$ weakly to V_* .

References

- [1] O.A. Alifanov, E.A. Artyukhin, S.V. Romyantsev. *Extremal methods for solving ill-posed problems*. M.: Nauka (1988)
- [2] O.Yu. Danilina. A nonlocal problem for the heat conduction equation with an integral condition (Russian). *Vestn. Samar. Gos. Tekhn. Univ., Ser. Fiz.-Mat. Nauki – J. Samara State Tech. Univ., Ser. Phys. Math. Sci.* **1**(14) (2007), 5–9.
- [3] A.D. Isgenderov. On variational statements of multidimensional inverse problems of mathematical physics (Russian). *DAN SSSR*. **274** (1984), no.3, 531-533.
- [4] A.D. Isgenderov, R.K. Tagiyev. Variational method solving the problem of identification of the coefficients of quasilinear parabolic equation. *The 7th International Conference “Inverse Problems: modelling and simulation” (IMPS-2014)*, May 26-31, Turkey. (2014) 31
- [5] N.I. Ivanchov. Boundary value problems for a parabolic equation with integral conditions (Russian). *Differential Equations*, **40**(4) (2004), 591–609.
- [6] S.I. Kabanikhin. *Inverse and ill-posed problems*. Novosibirsk: Siberian Scientific Publishing House (2009)
- [7] S.I. Kabanikhin, G.Dairbaeva. The inverse problem of finding the coefficient of the heat equation. *International conference Inverse ill-posed problems of mathematical physics, dedicated to the 75th anniversary of Academician M.M. Lavrentiev*, Russia, Novosibirsk, August 20-25, (2007) 1-5.
- [8] R.A. Kasumov. Difference approximation and regularization of the optimal control problem for a parabolic equation with controls in the coefficients and with a quality criterion along the boundary of the region. *Bulletin of the Baku University. Ser. Phys.-Math. Sciences.* (2016) no. 4. 61-76.
- [9] A.N.Kozhanov. On solvability of the boundary value problem with a nonlocal boundary condition for linear parabolic equations (Russian). *Vestn. Samar. Gos. Tekhn. Univ., Ser. Fiz.-Mat. Nauki – J. Samara State Tech. Univ., Ser. Phys. Math. Sci.* **30** (2004), 63–69.
- [10] O.A. Ladyzhenskaya. *Boundary value problems of mathematical physics*, Moscow, Nauka (1973), Russian.
- [11] F.V. Lubyshev. Difference approximations and regularization of optimal control problems for parabolic equations with controls in coefficients. *Zh. Comput. math. and mat. fiz.*, **35** (1995) no. 9 1313-1333
- [12] Sh.I. Maharramli. Inverse problem of control type on determining the higher coefficient of a one-dimensional parabolic equation. *Bulletin of the South Ural State University. Series: “Mathematics. Mechanics. Physics”*. **14** (2022). no. 1 35-41

- [13] A.A. Samarsky, V.B. Andreev. *Difference methods for elliptic equations*. M.: Nauka, (1976)
- [14] R.K. Tagiyev, Sh.I. Maharramli. On solvability of the initial-boundary problem for a one-dimensional linear parabolic equation with an integral boundary condition, *Vestn. Bakinskogo Univ., Ser. Fiz.-Mat. Nauki*, **2** (2019), 17-26, Russian
- [15] F.P. Vasil'ev, *Methods for solving extreme problems*. M.(1981).

Rafiq K. Tagiyev

Baku State University, Z. Khalilov 23, Baku, AZ1148, Azerbaijan

E-mail address: `r.tagiyev@list.ru`

Shahla I. Maharramli

Baku State University, Baku, Azerbaijan

E-mail address: `semedli.shehla@gmail.com`

Received: September 30, 2022; Revised: January 11, 2023; Accepted: January 17, 2023