

AVERAGING OPERATORS ON HERZ SEQUENCE SPACES WITH VARIABLE EXPONENTS

KWOK-PUN HO

Abstract. The main result of this paper establishes the boundedness of the averaging operators on the Herz sequence spaces with variable exponents. The Herz sequence spaces with variable exponents are extensions of the discrete Lebesgue spaces with variable exponent and the Herz spaces. We obtain our main result by extending the boundedness of the averaging operator on the discrete Lebesgue spaces with variable exponent to Herz sequence spaces with variable exponents.

1. Introduction

This paper gives the boundedness of the averaging operator on the Herz sequence spaces with variable exponents.

Recently, a number of function spaces on the Euclidean spaces had been discretized. For instance, we have the Morrey sequence spaces (the discrete Morrey spaces) introduced and studied in [5, 6, 7, 8]. We also have the discrete Lebesgue spaces with variable exponents [4, 22].

The Herz sequence spaces with variable exponents introduced in this paper are extensions of the discrete Lebesgue spaces with variable exponents. The discrete Lebesgue spaces with variable exponents are introduced in [22]. It is one of the important members of the modular space [21]. The embedding result for the discrete Lebesgue spaces with variable exponents was obtained in [23] and the boundedness of the averaging operator on the discrete Lebesgue spaces with variable exponents was established in [4]. The results obtained for the discrete Lebesgue spaces with variable exponents have applications on the studies of the Lebesgue spaces with variable exponents on Euclidean spaces [9, 24].

Moreover, the Herz sequence spaces with variable exponents are also the discrete analogue of the Herz space with variable exponents. The Herz spaces with variable exponents are extensions of the Lebesgue spaces with variable exponents [2, 3] and the Herz spaces [1, 10, 20, 25, 26, 27, 29].

This paper is organized as follows. Section 2 recalls the definitions of the discrete Lebesgue spaces with variable exponents and the boundedness of the averaging operators on the discrete Lebesgue spaces with variable exponents. The definition of the Herz sequence space with variable exponent is also given in

2000 *Mathematics Subject Classification.* 46E35, 46A45, 46B45, 26D15.

Key words and phrases. averaging operator, Herz spaces, sequence space, discrete Lebesgue spaces with variable exponents, boundedness.

this section. The main result on the boundedness of the averaging operators on the Herz sequence spaces with variable exponents is established in Section 3.

2. Preliminaries and Definitions

For any $a = \{a_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$ and $b = \{b_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$, we write $a \leq b$ if $a_n \leq b_n$, $\forall n \in \mathbb{Z}$. Define $|a| = \{|a_n|\}_{n \in \mathbb{Z}}$ and $\text{supp } a = \{n \in \mathbb{Z} : a_n \neq 0\}$.

Let $k \in \mathbb{N} \setminus \{0, 1\}$. For any $a = \{a_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$, the averaging operator $T_k a$ is defined as

$$(T_k a)_n = \frac{a_n + \cdots + a_{n+k-1}}{k}, \quad n \in \mathbb{Z}.$$

We now recall the definition of the discrete Lebesgue space with variable exponent from [4].

Definition 2.1. Let $\{p_n\}_{n \in \mathbb{Z}} \subset (1, \infty)$. The discrete Lebesgue space with variable exponent $l^{\{p_n\}}$ consists of all $a = \{a_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$ satisfying

$$\|a\|_{l^{\{p_n\}}} = \inf \left\{ \lambda > 0 : \sum_{n \in \mathbb{Z}} \left| \frac{a_n}{\lambda} \right|^{p_n} \leq 1 \right\} < \infty.$$

Whenever $p_n = p \in (1, \infty)$ for all $n \in \mathbb{Z}$, we see that $l^{\{p_n\}}$ becomes ℓ^p .

For the studies and applications of the discrete Lebesgue spaces with variable exponents, see [4, 9, 23]. The following is the condition that guarantees the boundedness of the averaging operator on $l^{\{p_n\}}$ from [4, Definition 3.5].

Definition 2.2. Let $\{p_n\}_{n \in \mathbb{Z}} \subset (1, \infty)$. We write $\{p_n\}_{n \in \mathbb{Z}} \in \mathcal{P}$ if there exist $\{\epsilon_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$ such that $p_n = p_0 + \epsilon_n$, $n \in \mathbb{Z}$ and $\sum_{n \in \mathbb{Z}} |\epsilon_n| < \infty$.

The following result is the boundedness of the averaging operator on $l^{\{p_n\}}$.

Proposition 2.1. Let $k \in \mathbb{N}$ and $\{p_n\}_{n \in \mathbb{Z}} \subset (1, \infty)$. If $\{p_n\}_{n \in \mathbb{Z}} \in \mathcal{P}$, then $T_k : l^{\{p_n\}} \rightarrow l^{\{p_n\}}$ is bounded.

For the proof of the above result, see [4, Theorem 3.6].

We now turn to the definition of the Herz sequence spaces with variable exponents. For any $i \in \mathbb{N} \setminus \{0\}$, write $D_i = \{n \in \mathbb{Z} : 2^i < n \leq 2^{i+1}\}$, $D_{-i} = \{n \in \mathbb{Z} : -2^{i+1} < n \leq -2^i\}$ and $D_0 = \{n \in \mathbb{Z} : |n| \leq 2\}$. For any $i \in \mathbb{Z} \setminus \{0\}$ and $a = \{a_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$, define $\chi_i a = \{(\chi_i a)_n\}_{n \in \mathbb{Z}}$ by

$$(\chi_i a)_n = \begin{cases} a_n, & n \in D_i, \\ 0, & n \notin D_i, \end{cases} \quad n \in \mathbb{Z}$$

and $\chi_0 a$ by

$$(\chi_0 a)_n = \begin{cases} a_n, & n \in D_0, \\ 0, & n \notin D_0, \end{cases} \quad n \in \mathbb{Z}.$$

It is easy to see that $\text{supp}(\chi_i a) = \text{supp } a \cap D_i$.

Definition 2.3. Let $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $\{p_n\}_{n \in \mathbb{Z}} \subset (1, \infty)$. The Herz sequence space with variable exponent $h_\theta^{\alpha, \{p_n\}}$ consists of all $a = \{a_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$ satisfying

$$\|a\|_{h_\theta^{\alpha, \{p_n\}}} = \left(\sum_{j \in \mathbb{Z}} (2^{j\alpha} \|\chi_j a\|_{l^{\{p_n\}}})^\theta \right)^{\frac{1}{\theta}} < \infty.$$

Let $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $p \in (1, \infty)$. The Herz sequence space $h_\theta^{\alpha, p}$ consists of all $a = \{a_n\}_{n \in \mathbb{Z}}$ satisfying

$$\|a\|_{h_\theta^{\alpha, p}} = \left(\sum_{j \in \mathbb{Z}} (2^{j\alpha} \|\chi_j a\|_{\ell^p})^\theta \right)^{\frac{1}{\theta}} < \infty.$$

Let $p \in (1, \infty)$. Whenever $p_n = p$ for all $n \in \mathbb{Z}$, the Herz sequence space with variable exponent $h_\theta^{\alpha, \{p_n\}}$ becomes the Herz sequence space $h_\theta^{\alpha, p}$.

The Herz sequence spaces with variable exponents are the extensions of the Herz spaces with variable exponents on Euclidean spaces in the discrete setting. For the studies of the Herz spaces with variable exponents on Euclidean spaces, see [11, 12, 13, 14, 15, 17, 18, 19, 28, 30].

3. Main result

This section establishes the main result of this paper. We extend the boundedness of the averaging operator to the Herz sequence spaces with variable exponents.

Theorem 3.1. *Let $k \in \mathbb{N}$, $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $\{p_n\}_{n \in \mathbb{Z}} \subset (1, \infty)$. If $\{p_n\}_{n \in \mathbb{Z}} \in \mathcal{P}$, then $T_k : h_\theta^{\alpha, \{p_n\}} \rightarrow h_\theta^{\alpha, \{p_n\}}$ is bounded.*

Proof. Let h be the smallest positive integer such that $k \leq 2^h$. Let $a = \{a_n\}_{n \in \mathbb{Z}} \in h_\theta^{\alpha, \{p_n\}}$.

For any $i \in \mathbb{N}$ with $2^{i-1} > k$, when $n \in D_i$, we have

$$n + k - 1 < 2^{i+1} + 2^{i-1} - 1 < 2^{i+2}.$$

Therefore,

$$\{n, \dots, n + k - 1\} \subset D_i \cup D_{i+1}.$$

Consequently,

$$|(\chi_i T_k a)_n| \leq \frac{|a_n| + \dots + |a_{n+k-1}|}{k} \leq T_k(|\chi_i a| + |\chi_{i+1} a|).$$

By applying the norm $\|\cdot\|_{l\{p_n\}}$ and, then multiplying $2^{i\alpha}$ on both sides of the above inequality, we obtain

$$2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}} \leq 2^{i\alpha} \|T_k(\chi_i a)\|_{l\{p_n\}} + 2^{i\alpha} \|T_k(\chi_{i+1} a)\|_{l\{p_n\}}.$$

Proposition 2.1 yields

$$2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}} \leq C 2^{i\alpha} (\|\chi_i a\|_{l\{p_n\}} + \|\chi_{i+1} a\|_{l\{p_n\}}). \quad (3.1)$$

When $n \in D_{-i}$, we have

$$n + k - 1 < -2^i + 2^{i-1} - 1 \leq -2^{i-1} - 1.$$

That is,

$$\{n, \dots, n + k - 1\} \subset D_i \cup D_{i-1}.$$

We find that

$$|(\chi_{-i} T_k a)_n| \leq \frac{|a_n| + \dots + |a_{n+k-1}|}{k} \leq (T_k(|\chi_{-i} a| + |\chi_{-i+1} a|))_n.$$

Therefore,

$$|\chi_i T_k a| \leq T_k(|\chi_{-i} a| + |\chi_{-i+1} a|).$$

Similar to (3.1), Proposition 2.1 gives

$$2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}} \leq C 2^{i\alpha} (\|\chi_{-i} a\|_{l\{p_n\}} + \|\chi_{-i+1} a\|_{l\{p_n\}}). \quad (3.2)$$

When i satisfies $k \geq 2^{i-1}$, we have $i \leq \frac{\ln k}{\ln 2} + 1$. For any $n \in D_{-i} \cup D_i$, we have

$$n + k - 1 \leq 2^{i+1} + 2^h - 1 \leq 2^{i+h+1}$$

because for any $b, c > 1$, we have $bc - c - b + 1 = (c-1)(b-1) > 0$. Consequently,

$$\{n, \dots, n+k-1\} \subset \bigcup_{j=-i}^{i+h} D_i$$

and

$$|(\chi_i T_k a)_n| \leq \frac{|a_n| + \dots + |a_{n+k-1}|}{k} \leq T_k \left(\sum_{j=-i}^{i+h} |\chi_j a| \right)$$

Therefore, Proposition 2.1 guarantees that

$$2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}} \leq C \left(\sum_{j=-i}^{i+h} 2^{i\alpha} \|\chi_j a\|_{l\{p_n\}} \right). \quad (3.3)$$

Write $m = \lfloor \frac{\ln k}{\ln 2} + 1 \rfloor$ where $[a]$ is the integral part of $a \in (0, \infty)$. As $\|\cdot\|_{\ell^\theta}$ is a quasi-norm, we have a constant $C > 0$ such that

$$\begin{aligned} \|\{2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}}\}_{i \in \mathbb{Z}}\|_{\ell^\theta} &\leq C \left(\left(\sum_{i=m}^{\infty} (2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}})^\theta \right)^{\frac{1}{\theta}} \right. \\ &\quad + \left(\sum_{i=-m+1}^{m-1} (2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}})^\theta \right)^{\frac{1}{\theta}} \\ &\quad \left. + \left(\sum_{i=-\infty}^{-m} (2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}})^\theta \right)^{\frac{1}{\theta}} \right) \\ &= I + II + III. \end{aligned}$$

According to (3.1), we have

$$I \leq C \left(\sum_{i=m}^{\infty} (2^{i\alpha} (\|\chi_i a\|_{l\{p_n\}} + \|\chi_{i+1} a\|_{l\{p_n\}}))^\theta \right)^{\frac{1}{\theta}} \leq C \|a\|_{h_\theta^{\alpha, \{p_n\}}} \quad (3.4)$$

for some $C > 0$ independent of a .

Similarly, (3.2) gives

$$\begin{aligned} III &\leq C \left(\sum_{i=-\infty}^{-m} (2^{i\alpha} (\|\chi_i a\|_{l\{p_n\}} + \|\chi_{i-1} a\|_{l\{p_n\}}))^\theta \right)^{\frac{1}{\theta}} \\ &\leq C \|a\|_{h_\theta^{\alpha, \{p_n\}}} \end{aligned} \quad (3.5)$$

for some $C > 0$ independent of a .

For II , for any $j \in \{-i, \dots, i+h\}$, since $0 \leq i \leq m$, we have $2^{(i-j)\alpha} \leq C$ because $-m-h \leq i-j \leq 2m$. Consequently, (3.3) yields

$$\begin{aligned} (2^{i\alpha} \|\chi_i T_k a\|_{l\{p_n\}})^\theta &\leq C \left(\sum_{j=-i}^{i+h} 2^{(i-j)\alpha} 2^{j\alpha} \|\chi_j a\|_{l\{p_n\}} \right)^\theta \\ &\leq C \left(\sum_{j=-i}^{i+h} (2^{j\alpha} \|\chi_j a\|_{l\{p_n\}})^\theta \right) \\ &\leq C \|a\|_{l\{p_n\}}^\theta \end{aligned}$$

for some $C > 0$ independent of i because there are at most $2i+h \leq m+h$ terms in the above summations. Therefore,

$$II \leq C \left(\sum_{i=-m+1}^{m-1} \sum_{j=-i}^{i+h} (2^{j\alpha} \|\chi_j a\|_{l\{p_n\}})^\theta \right)^{\frac{1}{\theta}} \leq C \|a\|_{h_\theta^{\alpha, \{p_n\}}} \quad (3.6)$$

for some $C > 0$ independent of a . Consequently, (3.4), (3.5) and (3.6) show that $T_k a \in h_\theta^{\alpha, \{p_n\}}$ and there is a constant $C > 0$ such that for any $a \in h_\theta^{\alpha, \{p_n\}}$, $\|T_k a\|_{h_\theta^{\alpha, \{p_n\}}} \leq C \|a\|_{h_\theta^{\alpha, \{p_n\}}}$. $\square$

Theorem 3.1 gives the following application on the discrete Herz spaces.

Corollary 3.1. *Let $k \in \mathbb{N}$, $\alpha \in \mathbb{R}$, $\theta \in (0, \infty)$ and $p \in (1, \infty)$. The averaging operator T_k is bounded on $h_\theta^{\alpha, p}$.*

Acknowledgments. The author thanks the reviewers for their valuable suggestions that improved the presentation of this paper.

References

- [1] J.M. Chen, J. Sun, Boundedness for Commutators of Singular Integral Operator With Weaker Smooth Kernel on Weighted Hardy and Herz Spaces, *Journal of Function Spaces*, **2024** (2024), Article ID 5596054.
- [2] D. Cruz-Uribe and A. Fiorenza, *Variable Lebesgue Spaces*, Birkhäuser, 2013.
- [3] L. Diening, P. Harjulehto, P. Hästö and M. Ružička, *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer, 2011.
- [4] D.E. Edmund and A. Nekvinda, Averaging operator on $l^{\{p_n\}}$ and $L^{p(\cdot)}$, *Math. Inequal. Appl.* **5** (2002), 235-246.
- [5] H. Gunawan, D. Hakim and M. Idris, On inclusion properties of discrete Morrey spaces, *Georgian Math. J.* **29** (2022), 37-44.
- [6] H. Gunawan, E. Kikianty and C. Schwanke, Discrete Morrey spaces and their inclusion properties, *Math. Nachr.* **291** (2018), 1283-1296.
- [7] D. Haroske and L. Skrzypczak, Morrey sequence spaces: Pittâ€™s theorem and compact embeddings, *Constr. Approx.* **51** (2020), 505-535.
- [8] D. Haroske and L. Skrzypczak, Generalised Morrey sequence spaces, *Adv. Oper. Theory* **10** (2025), Article Number: 61.
- [9] F. Hernández and C. Ruiz, Averaging and orthogonal operators on variable exponent spaces $L^{p(\cdot)}(\Omega)$, *J. Math. Anal. Appl.* **413** (2014), 139-153.
- [10] C.S. Herz, Lipschitz spaces and Bernsteinâ€™s theorem on absolutely convergent Fourier transforms, *J. Math. Mech* **18** (1968/69), 283-323.

- [11] K.-P. Ho, One-sided maximal operators on Herz spaces with variable exponents, *Math. Inequal. Appl.* **28** (2025), 189-198.
- [12] K.-P. Ho, Carleman-Knopp inequalities and geometric mean operators on Herz spaces with variable exponents, *Rocky Mount. J. Math.* (published online)
- [13] K.-P. Ho, Stein-Weiss inequalities on Herz spaces with variable exponent, *Z. Anal. Anwend.* (published online).
- [14] K.-P. Ho, One-sided operators on ball Banach function spaces, *J. Geom. Anal.* **35** (2025), Article Number: 276.
- [15] K.-P. Ho, General one-sided maximal operators on weighted one-sided Herz spaces, *J. Classical Anal.* (accepted)
- [16] K.-P. Ho, Herz spaces and integral operators on Morrey spaces. (preprint)
- [17] M. Izuki, Boundedness of commutators on Herz spaces with variable exponent, *Rend. Circ. Mat. Palermo* **59** (2010), 199-213.
- [18] M. Izuki, Y. Sawano, The Haar wavelet characterization of weighted Herz spaces and greediness of the Haar wavelet basis, *J. Math. Anal. Appl.* **362** (2010), 140-155.
- [19] M. Izuki, T. Noi and Y. Sawano, Extrapolation to two-weighted Herz spaces with three variable exponents, *Adv. Oper. Theory* **9** (2024), Article Number: 44.
- [20] S. Lu, D. Yang and G. Hu, *Herz type spaces and their applications*, Science Press, Beijing, 2008.
- [21] J. Musielak, *Orlicz Spaces and Modular Spaces Lecture Notes in Math.*, **1034**, Springer 1983.
- [22] H. Nakano, Modular sequence spaces, *Proc. Japan Acad.* **27** (1951), 411-415.
- [23] A. Nekvinda, Embeddings between discrete weighted Lebesgue spaces with variable exponents, *Math. Inequal. Appl.*, **10** (2007), 165-172.
- [24] A. Nekvinda, A note on maximal operator on $l^{\{p_n\}}$ and $L^{p(x)}(\mathbb{R})$, *J. Funct. Space* (2007).
- [25] M.A. Ragusa, Recent Advances in Homogeneous Herz Spaces and PDEs, AIP Conference Proceedings, **1168** (2009), 1492-1495.
- [26] M.A. Ragusa, Homogeneous Herz spaces and regularity results, *Nonlinear Anal.* **71** (2009), 1909-1914.
- [27] M.A. Ragusa, Parabolic Herz spaces and their applications, *Appl. Math. Lett.* **25** (2012), 1270-1273.
- [28] S. Samko, Variable exponent Herz spaces, *Mediterr. J. Math.* **10** (2013), 2007-2025.
- [29] M. Sultan and B. Sultan, A note on the boundedness of Marcinkiewicz integral operator on continual Herz-Morrey spaces, *Filomat* **39** (2025), 2017-2027.
- [30] T.-L. Yee and K.-P. Ho, Two weighted norm inequalities of potential type operator on Herz spaces, *Anal. Math.* **49** (2023), 1041-1052.

Kwok-Pun Ho

Department of Mathematics and Information Technology, The Education University of Hong Kong, 10, Lo Ping Road, Tai Po, Hong Kong, China

E-mail address: vkpho@eduhk.hk

Received: July 10, 2025; Revised: September 15, 2025; Accepted: October 2, 2025