

MULTI-SUBLINEAR ROUGH FRACTIONAL MAXIMAL OPERATOR ON PRODUCT MODIFIED MORREY SPACES

SABIR G. HASANOV

Abstract. We will study the boundedness of multi-sublinear fractional maximal operator $\mathcal{M}_{\Omega,\alpha,m}$ with rough kernels $\Omega \in L^s(\mathbb{S}^{mn-1})$, $1 < s \leq \infty$ on product modified Morrey spaces. We find for the operator $\mathcal{M}_{\Omega,\alpha,m}$ necessary and sufficient conditions on the parameters of the boundedness on product modified Morrey spaces $\tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$ to modified Morrey spaces $\tilde{L}^{q,\lambda}(\mathbb{R}^n)$.

1. Introduction

The classical Morrey spaces, introduced by Morrey [20] in 1938, have been studied intensively by various authors and together with weighted Lebesgue spaces play an important role in the theory of partial differential equations. They appeared to be quite useful in the study of local behavior of the solutions of elliptic differential equations and describe local regularity more precisely than Lebesgue spaces. See [4, 5, 6] for details. The boundedness of fractional integral operators on the classical Morrey spaces was studied by Adams [1], Chiarenza and Frasca *et al.* [3]. In [3], by establishing a pointwise estimate of fractional integrals in terms of the Hardy-Littlewood maximal function, they showed the boundedness of fractional integral operators on the Morrey spaces.

Let \mathbb{R}^n be the n -dimensional Euclidean space, and let $(\mathbb{R}^n)^m = \mathbb{R}^n \times \dots \times \mathbb{R}^n$ be the m -fold product space ($m \in \mathbb{N}$). For $x \in \mathbb{R}^n$ and $r > 0$, we denote by $B(x, r)$ the open ball centered at x of radius r , and by ${}^c B(x, r)$ denote its complement. Let $|B(x, r)|$ be the Lebesgue measure of the ball $B(x, r)$. Also for $\vec{x} = (x_1, \dots, x_m) \in \mathbb{R}^{mn}$ and $r > 0$, we denote by $B(\vec{x}, r)$ the open ball centered at $\vec{x} \in \mathbb{R}^{mn}$ of radius r . We denote by \vec{f} the m -tuple (f_1, f_2, \dots, f_m) , $\vec{y} = (y_1, \dots, y_m)$ and $d\vec{y} = dy_1 \cdots dy_m$.

Definition 1.1. Let $1 \leq p < \infty$, $0 \leq \lambda \leq \gamma$. We denote by $L_{p,\lambda}(\mathbb{R}^n)$ the Morrey space, and by $WL_{p,\lambda}(\mathbb{R}^n)$ the weak Morrey space, the set of locally integrable functions $f(x)$, $x \in \mathbb{R}^n$, with the finite norms

$$\|f\|_{L_{p,\lambda}} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))},$$

2010 *Mathematics Subject Classification.* Primary 42B20, 42B25, 42B35.

Key words and phrases. Multi-sublinear fractional maximal operator, Morrey spaces, modified Morrey spaces.

$$\|f\|_{WL_{p,\lambda}} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{WL_p(B(x,r))}$$

respectively.

Definition 1.2. Let $1 \leq p < \infty$, $0 \leq \lambda \leq \gamma$, $[r]_1 = \min\{1, r\}$. We denote by $\tilde{L}_{p,\lambda}(\mathbb{R}^n)$ the modified Morrey space, and by $W\tilde{L}_{p,\lambda}(\mathbb{R}^n)$ the weak modified Morrey space, the set of locally integrable functions $f(x)$, $x \in \mathbb{R}^n$, with the finite norms

$$\|f\|_{\tilde{L}_{p,\lambda}} = \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))},$$

$$\|f\|_{W\tilde{L}_{p,\lambda}} = \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{p}} \|f\|_{WL_p(B(x,r))}$$

respectively.

Note that

$$\tilde{L}_{p,0}(\mathbb{R}^n) = L_{p,0}(\mathbb{R}^n) = L_p(\mathbb{R}^n),$$

$$\tilde{L}_{p,\lambda}(\mathbb{R}^n) \subset_{\gamma} L_{p,\lambda}(\mathbb{R}^n) \cap L_p(\mathbb{R}^n) \quad \text{and} \quad \max\{\|f\|_{L_{p,\lambda}}, \|f\|_{L_p}\} \leq \|f\|_{\tilde{L}_{p,\lambda}}$$

and if $\lambda < 0$ or $\lambda > n$, then $L_{p,\lambda}(\mathbb{R}^n) = \tilde{L}_{p,\lambda}(\mathbb{R}^n) = \Theta$, where Θ is the set of all functions equivalent to 0 on \mathbb{R}^n .

The multilinear theory has been well developed in the past twenty years. In 1992, Grafakos [9] first study the following multilinear integrals, defined by

$$I_{\alpha}^m(\vec{f})(x) = \int_{\mathbb{R}^n} \frac{1}{|y|^{n-\alpha}} f_1(x - \theta_1 y) \dots f_m(x - \theta_m y) dy,$$

where $\theta_i (i = 1, \dots, m)$ are fixed distinct and nonzero real numbers and $0 < \alpha < n$. When $m = 1$ and $\Omega \equiv 1$, if let $\theta_1 = 1$, $I_{\Omega,\alpha,m}$ will be the Riesz potential operator I_{α} [22] given by

$$I_{\alpha} f(x) = \int_{\mathbb{R}^n} \frac{f(x-y)}{|y|^{n-\alpha}} dy.$$

Grafakos proved that the operator I_{α}^m is bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$ with $0 < 1/q = 1/p_1 + \dots + 1/p_m - \alpha/n < 1$, which can be regarded as an extension result for the classical fractional integral on Lebesgue spaces. In [11, 12, 15] was proved a certain O'Neil type inequality for dilated multi-linear convolution operators, including permutations of functions. This inequality was used to extend Grafakos result [9] to more general multi-linear operators of potential type and the relevant maximal operators.

In 1999, Kenig and Stein [19] studied the following multilinear fractional integral,

$$I_{\alpha,m}(\vec{f})(x) = \int_{(\mathbb{R}^n)^m} \frac{f_1(y_1) \dots f_m(y_m)}{|(x-y_1, \dots, x-y_m)|^{nm-\alpha}} dy_1 dy_2 \dots dy_m,$$

and showed that $I_{\alpha,m}$ is bounded from product $L_{p_1}(\mathbb{R}^n) \times L_{p_2}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_q(\mathbb{R}^n)$ with $1/q = 1/p_1 + \dots + 1/p_m - \alpha/n > 0$ for each $p_i > 1 (i = 1, \dots, m)$. If some $p_i = 1$, then $I_{\alpha,m}$ is bounded $L_{p_1}(\mathbb{R}^n) \times L_{p_2}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_{q,\infty}(\mathbb{R}^n)$. Obviously, the multilinear fractional integral $I_{\alpha,m}$ is a natural generalization of the classical fractional integral $I_{\alpha} \equiv I_{\alpha,1}$.

Let $1 < s \leq \infty$, $\Omega \in L^s(\mathbb{S}^{mn-1})$ be a homogeneous function of degree zero on \mathbb{R}^{mn} . The multi-sublinear fractional maximal operator $\mathcal{M}_{\alpha,m}$ with rough kernels Ω is defined by

$$\mathcal{M}_{\Omega,\alpha,m}(\vec{f})(x) = \sup_{r>0} \frac{1}{r^{nm-\alpha}} \int_{B(\vec{y},r)} |\Omega(\vec{y})| \prod_{j=1}^m |f_j(x - y_j)| d\vec{y}, \quad 0 \leq \alpha < nm.$$

If $m = 1$, then $M_{\Omega,\alpha} \equiv \mathcal{M}_{\Omega,\alpha,1}$ is the fractional maximal operator with rough kernel Ω . When $m = 1$ and $\Omega \equiv 1$, then $M_\alpha \equiv \mathcal{M}_{1,\alpha,1}$ is the classical fractional maximal operator.

Note that, in [13, 14] Guliyev, Ismayilova was study the boundedness of multi-sublinear fractional maximal operator, multilinear fractional integral operator and multilinear singular integral operators on product generalized Morrey spaces.

In this work, we prove the boundedness of the multi-sublinear fractional maximal operator with rough kernels $\mathcal{M}_{\Omega,\alpha,m}$ from product modified Morrey space $\tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$ to modified Morrey space $\tilde{L}^{q,\lambda}(\mathbb{R}^n)$, if $p > s'$, $1 < p_1, \dots, p_m < \infty$, $1/p = 1/p_1 + \dots + 1/p_m$, $\alpha/n \leq 1/p - 1/q \leq \alpha/(n - \lambda)$ and from the space $\tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$ to the weak space $W\tilde{L}^{q,\lambda}(\mathbb{R}^n)$, if $p = s'$, $1 \leq p_1, \dots, p_m < \infty$, $1/p = 1/p_1 + \dots + 1/p_m$, $\alpha/n \leq 1/s' - 1/q \leq \alpha/(n - \lambda)$ and at least one exponent p_i , $1 \leq i \leq m$ equals one.

Throughout this paper, we assume the letter C always remains to denote a positive constant that may vary at each occurrence but is independent of the essential variables.

2. Multi-sublinear fractional maximal operator $\mathcal{M}_{\Omega,\alpha,m}$ on product modified Morrey spaces

In this part, we investigate the boundedness of multi-sublinear fractional maximal operator $\mathcal{M}_{\Omega,\alpha,m}$ on product modified Morrey spaces.

Spanne and Adams obtained two remarkable results on Morrey spaces (see Definition 1.1 of the Morrey spaces in Section 1) for I_α . Their results can be summarized as follows.

Theorem 2.1. [16, 21] (*Spanne, but published by Peetre*) *Let $0 < \alpha < n$, $0 \leq \lambda < n - \alpha p$, $1/q = 1/p - \alpha/n$ and $\mu/q = \lambda/p$. Then for $p > 1$, the operators M_α and I_α are bounded from $L^{p,\lambda}(\mathbb{R}^n)$ to $L^{q,\mu}(\mathbb{R}^n)$ and for $p = 1$, I_α is bounded from $L^{1,\lambda}(\mathbb{R}^n)$ to $WL^{q,\mu}(\mathbb{R}^n)$.*

Theorem 2.2. [1, 10] *Let $0 < \alpha < n$, $1 \leq p < n/\alpha$, $0 \leq \lambda < n/\alpha p$.*

- (i) *If $p > 1$, then condition $1/p - 1/q = \alpha/(n - \lambda)$ is necessary and sufficient for the boundedness of the operator M_α from $L^{p,\lambda}(\mathbb{R}^n)$ to $L^{q,\lambda}(\mathbb{R}^n)$.*
- (ii) *If $p = 1$, then condition $1 - 1/q = \alpha/(n - \lambda)$ is necessary and sufficient for the boundedness of the operator M_α from $L^{1,\lambda}(\mathbb{R}^n)$ to $WL^{q,\lambda}(\mathbb{R}^n)$.*
- (iii) *If $p = \frac{n-\lambda}{\alpha}$, then the operator M_α is bounded from $L^{p,\lambda}(\mathbb{R}^n)$ to $L^\infty(\mathbb{R}^n)$.*

If $\lambda = 0$, then the statement of Theorems 2.1 and 2.2 reduces to the well known Hardy-Littlewood-Sobolev inequality.

On the other hand, in 2011, Guliyev, Hasanov and Zeren [10] found this inequality in modified Morrey spaces (see Definition 1.2 in Section 1) is also valid and proved that

Theorem 2.3. [10] *Let $0 < \alpha < n$, $1 \leq p < n/\alpha$, $0 \leq \lambda < n/\alpha p$.*

- (i) *If $p > 1$, then condition $\alpha/n \leq 1/p - 1/q \leq \alpha/(n - \lambda)$ is necessary and sufficient for the boundedness of the operator M_α from $\tilde{L}^{p,\lambda}(\mathbb{R}^n)$ to $\tilde{L}^{q,\lambda}(\mathbb{R}^n)$.*
- (ii) *If $p = 1$, then condition $\alpha/n \leq 1 - 1/q \leq \alpha/(n - \lambda)$ is necessary and sufficient for the boundedness of the operator M_α from $\tilde{L}^{1,\lambda}(\mathbb{R}^n)$ to $W\tilde{L}^{q,\lambda}(\mathbb{R}^n)$.*
- (iii) *If $\frac{n-\lambda}{\alpha} \leq p \leq \frac{n}{\alpha}$, then the operator M_α is bounded from $\tilde{L}^{p,\lambda}(\mathbb{R}^n)$ to $L^\infty(\mathbb{R}^n)$.*

When $m \geq 2$ and $\Omega \in L^s(\mathbb{S}^{mn-1})$, in [17] was find out $\mathcal{M}_{\Omega,m}$ also have the same properties by providing the following multi-version result of the Guliyev, Hasanov and Zeren [10].

Theorem 2.4. [17] *Let $1 < s \leq \infty$, $\Omega \in L^s(\mathbb{S}^{mn-1})$ be a homogeneous function of degree zero on \mathbb{R}^{mn} , p be the harmonic mean of $p_1, \dots, p_m > 1$ and*

$$\frac{\lambda}{p} = \sum_{j=1}^m \frac{\lambda_j}{p_j} \text{ for } 0 \leq \lambda_j < n. \quad (2.1)$$

- (i) *If $p > s'$, then the operator $\mathcal{M}_{\Omega,m}$ is bounded from product modified Morrey space $\tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$ to modified Morrey space $\tilde{L}^{p,\lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in \tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$*

$$\|\mathcal{M}_{\Omega,m}\mathbf{f}\|_{\tilde{L}^{p,\lambda}} \leq C \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j,\lambda_j}}.$$

- (ii) *If $p = s'$, then the operator $\mathcal{M}_{\Omega,m}$ is bounded from product modified Morrey space $\tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$ to weak modified Morrey space $W\tilde{L}^{p,\lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in \tilde{L}^{p_1,\lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m,\lambda_m}(\mathbb{R}^n)$*

$$\|\mathcal{M}_{\Omega,m}\mathbf{f}\|_{W\tilde{L}^{p,\lambda}} \leq C \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j,\lambda_j}}.$$

The following lemma was proved in [17].

Lemma 2.1. *Let $0 < \alpha < mn$, $1 \leq s' < mn/\alpha$, $\Omega \in L^s(\mathbb{S}^{mn-1})$ be a homogeneous function of degree zero on \mathbb{R}^{mn} , p be the harmonic mean of $p_1, \dots, p_m > 1$ and $f \in L_{\text{loc}}^1(\mathbb{R}^n) \times \dots \times L_{\text{loc}}^1(\mathbb{R}^n)$. Then for any $x \in \mathbb{R}^n$*

$$\mathcal{M}_{\Omega,\alpha,m}\mathbf{f}(x) \leq C_0 \prod_{j=1}^m \left[M_{\frac{\alpha s'}{m}}(f_j^{s'})(x) \right]^{\frac{1}{s'}} \leq C_0 \prod_{j=1}^m \left[M_{\frac{\alpha s' p_j}{m p}}(f_j^{\frac{s' p_j}{p}})(x) \right]^{\frac{p}{s' p_j}}, \quad (2.2)$$

where $C_0 = \frac{\|\Omega\|_{L^s(\mathbb{S}^{mn-1})}}{(mn)^{\frac{1}{s}}}$.

When $m \geq 2$ and $\Omega \in L^s(\mathbb{S}^{mn-1})$, in [18] was proved that for $\mathcal{M}_{\Omega, \alpha, m}$ also have the same properties by providing the following multi-version of the Theorem 2.2.

Theorem 2.5. *Let $0 < \alpha < mn$, $1 \leq s' < \frac{mn}{\alpha}$ and $\Omega \in L^s(\mathbb{S}^{mn-1})$. Suppose*

$$\sum_{j=1}^m \frac{\lambda_j}{p_j} = \frac{\lambda}{p}, \quad \sum_{j=1}^m \frac{\lambda_j}{q_j} = \frac{\lambda}{q}.$$

- (i) *If $p > s'$, $\frac{1}{p_j} - \frac{1}{q_j} = \frac{\alpha}{m(n-\lambda_j)}$ and $0 \leq \lambda_j < n - \frac{\alpha p_j}{m}$, then the condition $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ is necessary and sufficient for the boundedness of the operator $\mathcal{M}_{\Omega, \alpha, m}$ from product Morrey space $L^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times L^{p_m, \lambda_m}(\mathbb{R}^n)$ to $L^{q, \lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in L^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times L^{p_m, \lambda_m}(\mathbb{R}^n)$*

$$\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{L^{q, \lambda}} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}}.$$

- (ii) *If $p = s'$, $\frac{1}{p_j} - \frac{1}{p_j q_j} = \frac{\alpha}{m(n-\lambda_j)}$ and $0 \leq \lambda_j < n - \frac{\alpha p_j}{m}$, then the condition $\frac{1}{s'} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ is necessary and sufficient for the boundedness of the operator $\mathcal{M}_{\Omega, \alpha, m}$ from product Morrey space $L^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times L^{p_m, \lambda_m}(\mathbb{R}^n)$ to the weak Morrey space $WL^{q, \lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in L^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times L^{p_m, \lambda_m}(\mathbb{R}^n)$*

$$\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{WL^{q, \lambda}} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}}.$$

- (iii) *If $p = \frac{mn-\lambda}{\alpha} \geq s'$, then the operator $\mathcal{M}_{\Omega, \alpha, m}$ is bounded from $L^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times L^{p_m, \lambda_m}(\mathbb{R}^n)$ to $L^\infty(\mathbb{R}^n)$.*

When $m \geq 2$ and $\Omega \in L^s(\mathbb{S}^{mn-1})$, we find out $\mathcal{M}_{\Omega, \alpha, m}$ also have the same properties by providing the following multi-version of the Theorem 2.3.

Theorem 2.6. *Let $0 < \alpha < mn$, $1 \leq s' < \frac{mn}{\alpha}$ and $\Omega \in L^s(\mathbb{S}^{mn-1})$. Suppose*

$$\sum_{j=1}^m \frac{\lambda_j}{p_j} = \frac{\lambda}{p}, \quad \sum_{j=1}^m \frac{\lambda_j}{q_j} = \frac{\lambda}{q}.$$

- (i) *If $p > s'$, $\frac{\alpha}{mn} \leq \frac{1}{p_j} - \frac{1}{q_j} \leq \frac{\alpha}{m(n-\lambda_j)}$ and $0 \leq \lambda_j < n - \frac{\alpha p_j}{m}$, then the condition $\frac{\alpha}{n} \leq \frac{1}{p} - \frac{1}{q} \leq \frac{\alpha}{n-\lambda}$ is necessary and sufficient for the boundedness of the operator $\mathcal{M}_{\Omega, \alpha, m}$ from product modified Morrey space $\tilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$ to modified Morrey space $\tilde{L}^{q, \lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in \tilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$*

$$\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\tilde{L}^{q, \lambda}} \leq C \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j, \lambda_j}}.$$

- (ii) *If $p = s'$, $\frac{\alpha}{mn} \leq \frac{1}{p_j} - \frac{1}{p_j q_j} \leq \frac{\alpha}{m(n-\lambda_j)}$ and $0 \leq \lambda_j < n - \frac{\alpha p_j}{m}$, then the condition $\frac{\alpha}{n} \leq \frac{1}{s'} - \frac{1}{q} \leq \frac{\alpha}{n-\lambda}$ is necessary and sufficient for the boundedness of the operator $\mathcal{M}_{\Omega, \alpha, m}$ from product modified Morrey space $\tilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times$*

$\dots \times \widetilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$ to the weak modified Morrey space $W\widetilde{L}^{q, \lambda}(\mathbb{R}^n)$. Moreover, there exists a positive constant C such that for all $\mathbf{f} \in \widetilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \widetilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$

$$\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{W\widetilde{L}^{q, \lambda}} \leq C \prod_{j=1}^m \|f_j\|_{\widetilde{L}^{p_j, \lambda_j}}.$$

(iii) If $s' \leq \frac{mn-\lambda}{\alpha} \leq p \leq \frac{mn}{\alpha}$, then the operator $\mathcal{M}_{\Omega, \alpha, m}$ is bounded from $\widetilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \widetilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$ to $L^\infty(\mathbb{R}^n)$.

Proof. (i) *Sufficiency.* Taking the q -th root of both sides, applying Theorem 2.3 with $p/s' > 1$, the fact $\frac{\alpha}{mn} \leq \frac{1}{p_j} - \frac{1}{q_j} \leq \frac{\alpha}{m(n-\lambda_j)}$ and $f_j^{s'} \in \widetilde{L}^{p_j/s', \lambda_j}$, we get

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\widetilde{L}^{q, \lambda}} &= \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{B(x, t)} |\mathcal{M}_{\Omega, \alpha, m} f(y)|^q dy \right)^{1/q} \\ &\leq C \prod_{j=1}^m \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^{\lambda_j}} \int_{B(x, t)} \left[M_{\frac{\alpha s'}{m}}(f_j^{s'})(y) \right]^{\frac{q_j}{s'}} dy \right)^{\frac{1}{q_j}} \\ &= C \prod_{j=1}^m \left\| M_{\frac{\alpha s'}{m}}(f_j^{s'}) \right\|_{\widetilde{L}^{q_j/s', \lambda_j}}^{1/s'} \leq C \prod_{j=1}^m \|f_j^{s'}\|_{\widetilde{L}^{p_j/s', \lambda_j}}^{1/s'} = C \prod_{j=1}^m \|f_j\|_{\widetilde{L}^{p_j, \lambda_j}}, \end{aligned}$$

which is the desired inequality.

Necessity. Suppose that $\mathcal{M}_{\Omega, \alpha, m}$ is bounded from $\widetilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \widetilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$ to $\widetilde{L}^{q, \lambda}(\mathbb{R}^n)$. Define $\mathbf{f}_\varepsilon(x) = (f_1(\varepsilon x), \dots, f_m(\varepsilon x))$ for $\varepsilon > 0$. Then it is easy to show that

$$\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon(y) = \varepsilon^{-\alpha} \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(\varepsilon y). \quad (2.3)$$

Let $[t]_{1,+} = \max\{1, t\}$. Then by (2.3) we get

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon\|_{\widetilde{L}^{q, \lambda}} &= \varepsilon^{-\alpha} \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{B(x, t)} |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(\varepsilon y)|^q dy \right)^{\frac{1}{q}} \\ &= \varepsilon^{-\alpha - \frac{n}{q}} \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{B(\varepsilon x, \varepsilon t)} |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y)|^q dy \right)^{\frac{1}{q}} \\ &= \varepsilon^{-\alpha - \frac{n}{q}} \sup_{t > 0} \left(\frac{[\varepsilon t]_1}{[t]_1} \right)^{\frac{\lambda}{q}} \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[\varepsilon t]_1^\lambda} \int_{B(\varepsilon x, \varepsilon t)} |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y)|^q dy \right)^{\frac{1}{q}} \\ &= \varepsilon^{-\alpha - \frac{n}{q}} [\varepsilon]_{1,+}^{\frac{\lambda}{q}} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\widetilde{L}^{q, \lambda}}. \end{aligned}$$

Since $\mathcal{M}_{\Omega, \alpha, m}$ is bounded from $\widetilde{L}^{p_1, \lambda_1} \times \dots \times \widetilde{L}^{p_m, \lambda_m}$ to $\widetilde{L}^{q, \lambda}$, we have

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\widetilde{L}^{q, \lambda}} &= \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon\|_{\widetilde{L}^{q, \lambda}} \\ &\leq C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \|f_j(\varepsilon \cdot)\|_{\widetilde{L}^{p_j, \lambda_j}} \\ &= C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^{\lambda_j}} \int_{B(x, t)} |f_j(\varepsilon y)|^{p_j} dy \right)^{\frac{1}{p_j}} \end{aligned}$$

$$\begin{aligned}
&= C\varepsilon^{\alpha+\frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^{\lambda_j}} \int_{B(\varepsilon x, \varepsilon t)} |f_j(y)|^{p_j} dy \right)^{\frac{1}{p_j}} \\
&= C\varepsilon^{\alpha+\frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} \sup_{t > 0} \left(\frac{[\varepsilon t]_1}{[t]_1} \right)^{\frac{\lambda_j}{p_j}} \\
&\quad \times \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{(\varepsilon[t]_1)^{\lambda_j}} \int_{B(\varepsilon x, \varepsilon t)} |f_j(y)|^{p_j} dy \right)^{\frac{1}{p_j}} \\
&= C\varepsilon^{\alpha+\frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} [\varepsilon]_{1,+}^{\frac{\lambda_j}{p_j}} \|f_j\|_{\tilde{L}^{p_j, \lambda_j}},
\end{aligned}$$

where C is independent of ε .

If $n/p < n/q + \alpha$, then for all $\mathbf{f} \in \tilde{L}^{p_1, \lambda_1} \times \dots \times \tilde{L}^{p_m, \lambda_m}$, we have $\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\tilde{L}^{q, \lambda}} = 0$ as $\varepsilon \rightarrow 0$.

If $(n - \lambda)/p > (n - \lambda)/q + \alpha$, then for all $\mathbf{f} \in \tilde{L}^{p_1, \lambda_1} \times \dots \times \tilde{L}^{p_m, \lambda_m}$, we have $\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{\tilde{L}^{q, \lambda}} = 0$ as $\varepsilon \rightarrow \infty$.

Therefore we get $\frac{n-\lambda}{p} - \frac{n-\lambda}{q} \leq \alpha \leq \frac{n}{p} - \frac{n}{q}$.

- (ii) *Sufficiency.* If $p = s'$, for any $\beta > 0$, let $\varepsilon_0 = \beta$, $\varepsilon_m = 1$ and $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{m-1} > 0$ be arbitrary which will be chosen later. From the pointwise estimate (2.2), we get

$$\begin{aligned}
&\{y \in B(x, t) : |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y)| > \beta\} \\
&\subset \bigcup_{j=1}^m \left\{ y \in B(x, t) : \left[M_{\frac{\alpha s' p_j}{m p}} (f_j^{\frac{s' p_j}{p}})(x) \right]^{\frac{p}{s' p_j}} > \frac{\varepsilon_{j-1}}{t^{(\lambda - \lambda_j)/p_j \varepsilon_j}} \right\} \\
&= \bigcup_{j=1}^m \left\{ y \in B(x, t) : \left[M_{\frac{\alpha p_j}{m}} (f_j^{p_j})(x) \right]^{\frac{1}{p_j}} > \frac{\varepsilon_{j-1}}{t^{(\lambda - \lambda_j)/p_j \varepsilon_j}} \right\}.
\end{aligned}$$

Let us now take $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{m-1} > 0$ such that

$$\left(\frac{\varepsilon_j}{\varepsilon_{j-1}} \right)^{p_j q_j} = \frac{\left[\prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}} \right]^q}{\beta^q \|f_j\|_{L^{p_j, \lambda_j}}^{p_j q_j}}, \quad j = 1, 2, \dots, m.$$

Then, applying Lemma 2.1 with $p/s' = 1$ and the fact $f_j^{p_j} \in \tilde{L}^{1, \lambda_j}$, $\frac{\alpha}{mn} \leq \frac{1}{p_j} - \frac{1}{p_j q_j} \leq \frac{\alpha}{m(n - \lambda_j)}$ we get

$$\begin{aligned}
&\left| \{y \in B(x, t) : |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y)| > \beta\} \right| \\
&\leq C \sum_{j=1}^m \left| \left\{ y \in B(x, t) : \left[M_{\frac{\alpha p_j}{m}} (f_j^{p_j})(x) \right]^{\frac{1}{p_j}} > \frac{\varepsilon_{j-1}}{[t]_1^{(\lambda - \lambda_j)/(p_j q_j)} \varepsilon_j} \right\} \right| \\
&\leq C \sum_{j=1}^m [t]_1^{\lambda_j} \left(\frac{[t]_1^{(\lambda - \lambda_j)/(p_j q_j)} \varepsilon_j}{\varepsilon_{j-1}} \right)^{p_j q_j} \|f_j^{p_j}\|_{\tilde{L}^{1, \lambda_j}}^{q_j} \\
&= C \sum_{j=1}^m [t]_1^{\lambda} \left(\frac{\varepsilon_j}{\varepsilon_{j-1}} \|f_j\|_{\tilde{L}^{p_j, \lambda_j}} \right)^{p_j q_j}
\end{aligned}$$

$$= C \sum_{j=1}^m [t]_1^\lambda \left(\frac{1}{\beta} \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j, \lambda_j}} \right)^q = C [t]_1^\lambda \left(\frac{1}{\beta} \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j, \lambda_j}} \right)^q.$$

Hence, we obtain the following inequality

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{W\tilde{L}^{p, \lambda}} &= \sup_{\beta > 0} \beta \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{t^\lambda} \left| \left\{ y \in B(x, t) : |\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y)| > \beta \right\} \right| \right)^{\frac{1}{p}} \\ &\leq C \prod_{j=1}^m \|f_j\|_{\tilde{L}^{p_j, \lambda_j}}. \end{aligned}$$

This is the conclusion (ii) of Theorem 2.6.

Necessity. Suppose that $\mathcal{M}_{\Omega, \alpha, m}$ is bounded from $\tilde{L}^{p_1, \lambda_1}(\mathbb{R}^n) \times \dots \times \tilde{L}^{p_m, \lambda_m}(\mathbb{R}^n)$ to $W\tilde{L}_{q, \lambda}(\mathbb{R}^n)$. From equality (2.3) we get

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon\|_{W\tilde{L}^{q, \lambda}} &= \sup_{\tau > 0} \tau \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{\{y \in B(x, t) : \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon(y) > \tau\}} dy \right)^{1/q} \\ &= \sup_{\tau > 0} \tau \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{\{y \in B(x, t) : \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(\varepsilon y) > \tau \varepsilon^\alpha\}} dy \right)^{1/q} \\ &= \varepsilon^{-\frac{n}{q}} \sup_{\tau > 0} \tau \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^\lambda} \int_{\{y \in B(\varepsilon x, \varepsilon t) : \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(y) > \tau \varepsilon^\alpha\}} dy \right)^{1/q} \\ &= \varepsilon^{-\alpha - \frac{n}{q}} \sup_{t > 0} \left(\frac{[\varepsilon t]_1}{[t]_1} \right)^{\frac{\lambda}{q}} \sup_{\tau > 0} \tau \varepsilon^\alpha \\ &\quad \times \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{(\varepsilon [t]_1)^\lambda} \int_{\{y \in B(x, \varepsilon t) : \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(\varepsilon y) > \tau \varepsilon^\alpha\}} dy \right)^{1/q} \\ &= \varepsilon^{-\alpha - \frac{n}{q}} [\varepsilon]_{1, +}^{\frac{\lambda}{q}} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{W\tilde{L}^{q, \lambda}}. \end{aligned}$$

By the boundedness of the operator $\mathcal{M}_{\Omega, \alpha, m}$ from $\tilde{L}^{p_1, \lambda_1} \times \dots \times \tilde{L}^{p_m, \lambda_m}$ to $W\tilde{L}^{q, \lambda}$, we have

$$\begin{aligned} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{W\tilde{L}^{q, \lambda}} &= \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1, +}^{-\frac{\lambda}{q}} \|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}_\varepsilon\|_{W\tilde{L}^{q, \lambda}} \\ &\leq C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1, +}^{-\frac{\lambda}{q}} \prod_{j=1}^m \|f_j(\varepsilon \cdot)\|_{\tilde{L}^{p_j, \lambda_j}} \\ &= C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1, +}^{-\frac{\lambda}{q}} \prod_{j=1}^m \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^{\lambda_j}} \int_{B(x, t)} |f_j(\varepsilon y)|^{p_j} dy \right)^{1/p_j} \\ &= C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1, +}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{[t]_1^{\lambda_j}} \int_{B(\varepsilon x, \varepsilon t)} |f_j(y)|^{p_j} dy \right)^{1/p_j} \\ &= C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1, +}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} \sup_{t > 0} \left(\frac{[\varepsilon t]_1}{[t]_1} \right)^{\frac{\lambda_j}{p_j}} \\ &\quad \times \sup_{x \in \mathbb{R}^n, t > 0} \left(\frac{1}{(\varepsilon [t]_1)^{\lambda_j}} \int_{B(\varepsilon x, \varepsilon t)} |f_j(y)|^{p_j} dy \right)^{1/p_j} \end{aligned}$$

$$= C \varepsilon^{\alpha + \frac{n}{q}} [\varepsilon]_{1,+}^{-\frac{\lambda}{q}} \prod_{j=1}^m \varepsilon^{-\frac{n}{p_j}} [\varepsilon]_{1,+}^{\frac{\lambda_j}{p_j}} \|f_j\|_{\tilde{L}^{p_j, \lambda_j}},$$

where C is independent of ε .

If $1/p < 1/q + \alpha/n$, then for all $\mathbf{f} \in \tilde{L}^{p_1, \lambda_1} \times \dots \times \tilde{L}^{p_m, \lambda_m}$, we have $\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{WL^q, \lambda} = 0$ as $\varepsilon \rightarrow 0$.

If $1/p > 1/q + \alpha/(n - \lambda)$, then for all $\mathbf{f} \in \tilde{L}^{p_1, \lambda_1} \times \dots \times \tilde{L}^{p_m, \lambda_m}$, we have $\|\mathcal{M}_{\Omega, \alpha, m} \mathbf{f}\|_{WL^q, \lambda} = 0$ as $\varepsilon \rightarrow \infty$.

Therefore we get $\alpha/n \leq 1/p - 1/q \leq \alpha/(n - \lambda)$.

(iii) Since $\Omega \in L^s(\mathbb{S}^{mn-1})$ with $s' \leq \frac{mn-\lambda}{\alpha} \leq p \leq \frac{mn}{\alpha}$, Hölder's inequality yields that

$$\begin{aligned} \mathcal{M}_{\Omega, \alpha, m} \mathbf{f}(x) &= \sup_{r>0} \frac{1}{r^{nm-\alpha}} \int_{B(\vec{y}, r)} |\Omega(\vec{y})| \prod_{j=1}^m |f_j(x - y_j)| d\vec{y} \\ &\leq \sup_{r>0} \frac{1}{r^{nm-\alpha}} \left(\int_{B(\vec{y}, r)} \prod_{j=1}^m |f_j(x - y_j)|^p d\vec{y} \right)^{\frac{1}{p}} \left(\int_{B(\vec{y}, r)} |\Omega(\vec{y})|^{p'} d\vec{y} \right)^{\frac{1}{p'}} \\ &\leq \sup_{r>0} \frac{1}{r^{nm-\alpha}} \left(\int_{B(\vec{y}, r)} \prod_{j=1}^m |f_j(x - y_j)|^p d\vec{y} \right)^{\frac{1}{p}} \\ &\quad \times \left(\int_{B(\vec{y}, r)} |\Omega(\vec{y})|^s d\vec{y} \right)^{\frac{1}{s}} |B(\vec{y}, r)|^{\frac{1}{p'} - \frac{1}{s}} \\ &\leq C \sup_{r>0} \frac{r^{nm(\frac{1}{p'} - \frac{1}{s})}}{r^{nm-\alpha}} \left(\int_{B(\vec{y}, r)} \prod_{j=1}^m |f_j(x - y_j)|^p d\vec{y} \right)^{\frac{1}{p}} \left(\int_{B(\vec{y}, r)} |\Omega(\vec{y})|^s d\vec{y} \right)^{\frac{1}{s}} \\ &= C \sup_{r>0} r^{\alpha - \frac{nm}{p} - \frac{nm}{s}} \prod_{j=1}^m \left(\int_{B(y, r)} |f_j(x - y_j)|^{p_j} d\vec{y} \right)^{\frac{1}{p_j}} \\ &\quad \times \left(\int_0^r \int_{\mathbb{S}^{mn-1}} |\Omega(\xi)|^s t^{mn-1} d\xi dt \right)^{\frac{1}{s}} \\ &= C \sup_{r>0} r^{\alpha - \frac{nm}{p}} \prod_{j=1}^m \left(\int_{B(y, r)} |f_j(x - y_j)|^{p_j} dy_j \right)^{\frac{1}{p_j}} \\ &\leq C \sup_{r>0} r^{\alpha - \frac{nm}{p}} \prod_{j=1}^m [r]_1^{\frac{\lambda_j}{p_j}} \prod_{j=1}^m \left(\frac{1}{[r]_1^{\lambda_j}} \int_{B(y, r)} |f_j(x - y_i)|^{p_j} dy_i \right)^{\frac{1}{p_j}} \\ &\leq C \sup_{r>0} r^{\alpha - \frac{nm}{p}} [r]_1^{\frac{\lambda}{p}} \prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}} \\ &\leq C \prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}} \max \left\{ \sup_{0 < r < 1} r^{\alpha - \frac{nm}{p}}, \sup_{r \geq 1} r^{\alpha - \frac{nm}{p} + \frac{\lambda}{p}} \right\} \leq C \prod_{j=1}^m \|f_j\|_{L^{p_j, \lambda_j}}. \end{aligned}$$

□

References

- [1] D. R. Adams, A note on Riesz potentials, *Duke Math.*, **42**, (1975), 765–778.
- [2] V. I. Burenkov, *Sobolev spaces on domains*, B.G. Teubner, Stuttgart-Leipzig, 1998.
- [3] F. Chiarenza, M. Frasca, Morrey spaces and Hardy-Littlewood maximal function, *Rend Mat.*, **7** (1987), 273–279.
- [4] F. Chiarenza, M. Frasca, P. Longo, Interior $W^{2,p}$ -estimates for nondivergence elliptic equations with discontinuous coefficients, *Ricerche Mat.*, **40** (1991), 149–168.
- [5] F. Chiarenza, M. Frasca, P. Longo, $W^{2,p}$ -solvability of Dirichlet problem for non-divergence elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.*, **336** (1993), 841–853.
- [6] G. Di Fazio, D. K. Palagachev, M. A. Ragusa, Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients, *J. Funct. Anal.*, **166**, no. 2 (1999), 179–196.
- [7] Y. Ding and S. Lu, The $L^{p_1} \times L^{p_2} \times \dots \times L^{p_k}$ boundedness for some rough operators, *J. Math. Anal. Appl.*, **203** (1996), 166–186.
- [8] H. Gunawan, I. Sihwaningrum, Multilinear maximal functions and fractional integrals on generalized Morrey spaces, *J. Math. Anal.*, **4**, no. 3 (2013), 23–32.
- [9] L. Grafakos, On multilinear fractional integrals, *Studia Math.*, **102** (1992), 49–56.
- [10] V. Guliyev, J. Hasanov and Y. Zeren, Necessary and sufficient conditions for the boundedness of the Riesz potential in modified Morrey spaces, *J. Math. Inequal.*, **5** (2011), 491–506.
- [11] V. S. Guliyev, Sh. A. Nazirova, A rearrangement estimate for the generalized multilinear fractional integrals, *Siberian Math. J.*, **48** (2007), 463–470.
- [12] V. S. Guliyev, Sh. A. Nazirova, O’Neil inequality for multilinear convolutions and some applications, *Integral Equations Operator Theory*, **60** (2008), 485–497.
- [13] V. S. Guliyev, A. F. Ismayilova, Multi-sublinear fractional maximal operator and multilinear fractional integral operators on generalized Morrey spaces, *Proceedings of the Institute of Mathematics and Mechanics*, **40**, no. 2, 2014, 22–33.
- [14] V. S. Guliyev, A. F. Ismayilova, Multi-sublinear maximal operator and multilinear singular integral operators on generalized Morrey spaces. *Proceedings of the Institute of Mathematics and Mechanics*, **40**, no. 2, 2014, 65–77.
- [15] V. S. Guliyev, I. Ekincioglu, Sh. A. Nazirova, The $L_{p_1 r_1} \times L_{p_2 r_2} \times \dots \times L_{p_k r_k}$ boundedness of rough multilinear fractional integral operators in the Lorentz spaces, *Journal of Inequalities and Applications* (2015), 2015:71.
- [16] V. Guliyev and P. Shukurov, On the boundedness of the fractional maximal operator, Riesz potential and their commutators in generalized Morrey spaces, *Oper. Theory Adv. Appl.*, **229** (2013), 175–199.
- [17] S. G. Hasanov, Multi-sublinear rough maximal operator on product Morrey and product modified Morrey spaces, *Journal of Contemporary Applied Mathematics*, **4**, no. 2 (2014), 57–65.
- [18] S. G. Hasanov, Multi-sublinear rough fractional maximal operator on product Morrey spaces, *Journal of Contemporary Applied Mathematics*, **4**, no. 2 (2014), 66–76.
- [19] C. E. Kenig, E. M. Stein, Multilinear estimates and fractional integration, *Math. Res. Lett.*, **6** (1999), 1–15.
- [20] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.*, **43** (1938), 126–166.
- [21] J. Peetre, On the theory of $L^{p,\lambda}$, *J. Funct. Anal.* **4** (1969), 71–87.
- [22] E. M. Stein, *Singular integrals and differentiability properties of functions*. Princeton Univ. Press, Princeton, NJ, 1970.

Gandja State University, Gandja, Azerbaijan

E-mail address: sabirhasanov@gmail.com

Received: March 17, 2015; Accepted: May 15, 2015